

Holographic Techni-dilaton *

Kazumoto Haba,^{1,†} Shinya Matsuzaki,^{2,‡} and Koichi Yamawaki^{3,§}

¹*Department of Physics, Nagoya University, Nagoya, 464-8602, Japan.*

²*Department of Physics, Pusan National University, Busan 609-735, Korea.*

³*Kobayashi-Maskawa Institute for the Origin of Particles and the Universe (KMI)
Nagoya University, Nagoya 464-8602, Japan.*

(Dated: August 6, 2010)

Techni-dilaton, a pseudo-Nambu-Goldstone boson of scale symmetry, was predicted long ago in the Scale-invariant/Walking/Conformal Technicolor (SWC-TC) as a remnant of the (approximate) scale symmetry associated with the conformal fixed point, based on the conformal gauge dynamics of ladder Schwinger-Dyson (SD) equation with non-running coupling. We study the techni-dilaton as a flavor-singlet bound state of techni-fermions by including the techni-gluon condensate (tGC) effect into the previous (bottom-up) holographic approach to the SWC-TC, a deformation of the holographic QCD with $\gamma_m \simeq 0$ by large anomalous dimension $\gamma_m \simeq 1$. With including a bulk scalar field corresponding to the gluon condensate, we first improve the Operator Product Expansion of the current correlators so as to reproduce gluonic $1/Q^4$ term both in QCD and SWC-TC. We find in QCD about 10% (negative) contribution of gluon condensate to the ρ meson mass. We also calculate the oblique electroweak S -parameter in the presence of the effect of the tGC and find that for the fixed value of S the tGC effects dramatically reduce the flavor-singlet scalar (techni-dilaton) mass M_{TD} (in the unit of F_π), while the vector and axial-vector masses M_ρ and M_{a_1} are rather insensitive to the tGC, where F_π is the decay constant of the techni-pion. If we use the range of values of tGC implied by the ladder SD analysis of the non-perturbative scale anomaly in the large N_f QCD near the conformal window, the phenomenological constraint $S \simeq 0.1$ predicts the techni-dilaton mass $M_{TD} \sim 600$ GeV which is within reach of LHC discovery.

PACS numbers:

I. INTRODUCTION

The origin of mass is the most urgent issue of the particle physics today and is to be resolved at the LHC experiments. In the standard model, all masses are attributed to a single parameter of the vacuum expectation value (VEV) of the hypothetical elementary particle, the Higgs boson, which is simply transferred from the input mass parameter μ tuned as tachyonic ($\mu^2 < 0$) in an ad hoc manner. As such the standard model does not explain the origin of mass.

Technicolor (TC) [2] is an attractive idea to account for the origin of mass without introducing Higgs boson and tachyonic mass parameter in analogy with QCD: The mass arises dynamically from the condensate of the fermion-antifermion pair like Cooper pair in the Bardeen-Cooper-Schrieffer theory of the superconductor, picking up the intrinsic mass scale Λ_{TC} generated by the scale anomaly [3] through quantum effects in the gauge theory which is scale-invariant at classical level for massless flavors. Actually in QCD, the running of the coupling constant $\alpha(\mu)$ implies existence of the intrinsic mass scale Λ_{QCD} . The original version of TC, just a simple scale-up of QCD, however, is plagued by the notorious problems: Excessive flavor-changing neutral currents (FCNCs), and excessive oblique corrections of $\mathcal{O}(1)$ to the Peskin-Takeuchi S parameter [4] compared with the typical experimental bound about 0.1.

The FCNC problem was resolved long time ago by the scale-invariant/walking/conformal TC (SWC-TC) [5–7][8] initially dubbed as “scale-invariant TC” with the prediction of a “techni-dilaton”, a pseudo Nambu-Goldstone boson of the spontaneous breaking of the (approximate) scale invariance [5]. The theory was based on the strong coupling solution of Maskawa-Nakajima [9] for the ladder Schwinger-Dyson (SD) equation with *non-running* (conformal) gauge coupling $\alpha > \alpha_{cr} = \mathcal{O}(1)$. It was found [5] (for reviews, see [10]) that the solution implies a large anomalous dimension $\gamma_m \simeq 1$ of the techni-fermion condensate operator $\bar{T}T$ at $\alpha \simeq \alpha_{cr}$, namely the enhanced condensate

* Short report of this paper was given at [1].

[†]Present address: MRI Research Associates Inc., Tokyo 101-0047, Japan. E-mail: kazumohaba@gmail.com

[‡]synya@pusan.ac.kr

[§]yamawaki@kmi.nagoya-u.ac.jp

$\frac{\langle \bar{T}T \rangle_\Lambda}{\langle \bar{T}T \rangle_m} = \left(\frac{\Lambda}{m}\right)^{\gamma_m} \gtrsim \frac{\Lambda}{m}$ to avoid the FCNC problem, where m is the dynamical mass of techni-fermion on the order of weak scale $\simeq 250$ GeV and the cutoff Λ is usually identified with the scale of Extended Technicolor (ETC) [11], $\Lambda = \Lambda_{\text{ETC}} (> 10^3 m)$. In contrast to a folklore that TC is a “higgsless model”, a salient feature of SWC-TC is the conformality which manifests itself by the appearance of a composite Higgs boson (“conformal Higgs”) as the techni-dilaton proposed initially in reference [5]. It is this (approximate) conformal symmetry that is responsible for the naturalness of the SWC-TC to guarantee the large hierarchy $m \ll \Lambda$ in such a way that the coupling is almost non-running (conformal) over the wide energy range $m < \mu < \Lambda$. Moreover, there also exists a possibility [12, 13] that the S parameter can be reduced in the case of SWC-TC.

An explicit gauge dynamics [14, 15] of such an SWC-TC is based on the Caswell-Banks-Zaks infrared fixed point [16, 17] (CBZ-IRFP) $\alpha_* = \alpha_*(N_c, N_f)$ which appears in the two-loop beta function of the “large N_f QCD”, QCD with the number of massless flavors $N_f (< 11N_c/2)$ larger than a certain value $N_f^* (> N_c)$. Note that $\alpha_* \searrow 0$ when $N_f \nearrow 11N_c/2$, and hence there exists a certain region ($N_f^* < N_f^{\text{cr}} < N_f < 11N_c/2$) (“conformal window”) such that $\alpha_* < \alpha_{\text{cr}}$, where α_{cr} is the critical coupling for the spontaneous chiral symmetry breaking and hence the chiral symmetry gets restored in this region. Here α_{cr} may be evaluated as $\alpha_{\text{cr}} = \pi/3C_2(F)$ in the ladder approximation [18], in which case $\alpha_* = \alpha_*(N_c, N_f^{\text{cr}}) = \alpha_{\text{cr}}$ determines N_f^{cr} as $N_f^{\text{cr}} \simeq 4N_c$ [14] ^{#1} ^{#2}. Related to the conformal symmetry, this phase transition (“conformal phase transition” [15]) has unusual nature that the order parameter changes continuously but the spectrum does discontinuously at the phase transition point $\alpha_* = \alpha_{\text{cr}}$ when we change α_* (or N_f/N_c) continuously.

When we set α_* slightly larger than α_{cr} (slightly outside of the conformal window), the walking coupling $\alpha(\mu) (< \alpha_*)$ becomes larger than the critical coupling in the wide infrared region, we have a condensate or the dynamical mass of the techni-fermion m , which is much smaller than the intrinsic scale of the theory $\Lambda_{\text{TC}} (> m)$. Such an intrinsic scale Λ_{TC} is quantum mechanically induced by the scale anomaly as an analogue of Λ_{QCD} in QCD and the theory behaves as ordinary asymptotically free theory as QCD for $\mu > \Lambda_{\text{TC}}$ (Region I in Fig. 1). Although the CBZ-IRFP α_* actually disappears (then would-be IRFP) at the scale $\mu \sim m$ where the fermions have acquired the mass m and get decoupled from the beta function for $\mu < m$ (Region III in Fig. 1), the coupling is still walking due to the remnant of the CBZ-IRFP conformality in a wide region $m < \mu < \Lambda_{\text{TC}}$ (Region II in Fig. 1). Then the theory acts like an SWC-TC: Λ_{TC} plays a role of cutoff Λ identified with the ETC scale: $\Lambda_{\text{TC}} = \Lambda = \Lambda_{\text{ETC}}$. It develops a large anomalous dimension $\gamma_m \simeq 1$ for Region II to solve the FCNC problem [14, 15].




FIG. 1: The beta function and $\alpha(\mu)$ for SWC-TC.

Existence of these two largely separated scales, m and Λ_{TC} such that $m \ll \Lambda_{\text{TC}}$, is the most important feature of SWC-TC, in sharp contrast to the ordinary QCD with small number of flavors (in the chiral limit) where all the mass parameters like dynamical mass of quarks are of order of the single scale parameter of the theory Λ_{QCD} , $m \sim \Lambda_{\text{QCD}}$. The intrinsic scale Λ_{TC} is related with the scale anomaly corresponding to the *perturbative* running effects of the coupling, with the ordinary beta function $\beta(\alpha)$ in the Region I, in the same sense as in QCD [3].

$$\langle \partial^\mu D_\mu \rangle = \langle \theta_\mu^\mu \rangle = 4 \langle \theta_0^0 \rangle = \frac{\beta(\alpha)}{4\alpha^2} \langle \alpha G_{\mu\nu}^2 \rangle = \mathcal{O}(\Lambda_{\text{TC}}^4), \quad (1.1)$$

which implies that all the techni-gluon balls have mass of $\mathcal{O}(\Lambda_{\text{TC}})$. On the other hand, the scale m is related with totally different scale anomaly due to the dynamical generation of m which does exist even in the idealized case

^{#1} In the case of $N_c = 3$, this value $N_f^{\text{cr}} \simeq 4N_c = 12$ is somewhat different from the lattice value [19] $6 < N_f^{\text{cr}} < 7$, but is consistent with more recent lattice results [20].

^{#2} There is another possibility for the SWC-TC with much less N_f based on the higher TC representation [21], although explicit ETC model building would be somewhat involved.

with non-running coupling $\alpha(\mu) \equiv \alpha(> \alpha_{\text{cr}})$ such as the Maskawa-Nakajima solution [9], as was discussed some time ago [22]. Such an idealized case well simulates the dynamics of Region II [14, 15], with anomalous dimension $\gamma_m \simeq 1$ and $m \ll \Lambda_{\text{TC}}$ in the numerical calculations [23] with the walking coupling constant in the most of the Region II being slightly larger than α_{cr} , $\alpha_* > \alpha(\mu) > \alpha_{\text{cr}}$. The coupling $\alpha \equiv \alpha_*$ in the “idealized Region II” actually runs non-perturbatively according to the essential-singularity scaling (Miransky scaling [24]) of mass generation, with the *non-perturbative* beta function $\beta_{\text{NP}}(\alpha)$. Then the *non-perturbative* scale anomaly reads [15]

$$\langle \partial^\mu D_\mu \rangle = \langle \theta_\mu^\mu \rangle = 4\langle \theta_0^0 \rangle = \frac{\beta_{\text{NP}}(\alpha)}{4\alpha^2} \langle \alpha G_{\mu\nu}^2 \rangle = \mathcal{O}(m^4), \quad (1.2)$$

which vanishes when we approaches the conformal window from the broken phase $\alpha_* \searrow \alpha_{\text{cr}}$ ($m \rightarrow 0$) where α_* is the would-be CBZ-IRFP (See Eqs.(4.3) and (4.4)). All the techni-fermion bound states have mass of order of m , while there are no light bound states in the symmetric phase (conformal window) $\alpha_* < \alpha_{\text{cr}}$, a characteristic feature of the conformal phase transition [15]. The techni-dilaton is associated with the latter scale anomaly and should have mass on order of $m(\ll \Lambda_{\text{TC}})$.

More concretely, the mass of techni-dilaton or scalar bound state in the SWC-TC was estimated in various methods: The first method was based on the assumption of partially conserved dilatation current (PCDC) [25] combined with the ladder SD equation for the gauged Nambu-Jona-Lasinio model which well simulates [14, 15] the conformal phase transition in the large N_f QCD. The result indicates

$$M_{\text{TD}} \simeq \sqrt{2}m, \quad (1.3)$$

which coincides with other methods in the ladder SD equation without use of the PCDC [26]. Also a straightforward calculation [23, 27] of scalar bound state mass as well as the S parameter [13] was made in the vicinity of the CBZ-IRFP in the large N_f QCD, based on the coupled use of the SD equation and the Bethe-Salpeter (BS) equation in the ladder approximation:

$$M_{\text{TD}} \sim 1.5m (< M_\rho, M_{a_1}), \quad (1.4)$$

where M_ρ, M_{a_1} are masses of (techni-) ρ and (techni-) a_1 mesons, respectively, which is consistent with Eq.(1.3) in contrast to the ordinary QCD where the scalar mass is larger than those of the vector mesons (“higgsless”) within the same framework of ladder SD/BS equation approach.

The SWC-TC, however, has a calculability problem, since its non-perturbative dynamics is not QCD-like at all, and hence no simple scaling of QCD results would be available. The best thing we could do so far has been a straightforward calculation based on the SD equation and (inhomogeneous) BS equation in the ladder approximation [13], which is however not a systematic approximation and is not very reliable in the quantitative sense.

Of a late fashion, based on the so-called AdS/CFT (anti-de-Sitter space/conformal field theory) correspondence, a duality of the string in the anti-de Sitter space background-conformal field theory [28], holography gives us a new method which may resolve the calculability problem of strongly coupled gauge theories [29]: Use of the holographic correspondence enables us to calculate Green functions in a four-dimensional strongly coupled theory from a five-dimensional weakly coupled theory. For instance, QCD can be reformulated based on the holographic correspondence either in the bottom-up approach [30, 31] or in the top-down approach [32]. In both approaches we end up with the five-dimensional gauge theory for the flavor symmetry, whose infinite tower of Kaluza-Klein (KK) modes describe nicely a set of the massive vector/axial-vector mesons as the gauge bosons of Hidden Local Symmetries (HLSs) [33, 34], or equivalently as the Moose [35]. Although a holographic description is valid only for large N_c limit, several observables of QCD have been reproduced within 30 % errors in both approaches. Moreover, through the high-energy behavior of current correlators in operator product expansion (OPE), some consistency with the QCD has been confirmed in the bottom-up approach.

Recently the S parameter in the SWC-TC was calculated [36–39] as an application of the above technique of bottom-up holographic QCD (hard-wall model) [30, 31] to the holographic SWC-TC. In the previous work [39], based on the holographic correspondence in the bottom-up approach, we calculated the S parameter in the SWC-TC, treating the anomalous dimension γ_m as a free parameter as $0 \lesssim \gamma_m \lesssim 1$, varying continuously from the QCD monitor value $\gamma_m \simeq 0$ through the one of the SWC-TC $\gamma_m \simeq 1$. We obtained S as an explicit function of F_π/M_ρ in entire region, which turns out to be a positive and monotonically increasing function such that S continuously goes to zero when $F_\pi/M_\rho \rightarrow 0$, where F_π and M_ρ are the (techni-)pion decay constant and the (techni-) ρ meson mass, respectively.

In this paper, we extend the previous paper [39] on the hard-wall-type bottom-up holographic SWC-TC by including effects of (techni-) gluon condensation, Γ , through the bulk flavor/chiral-singlet scalar field Φ_X , in addition to the conventional bulk scalar field Φ dual to the chiral condensate. For definition of Γ , see text. The techni-dilaton, a flavor-singlet scalar bound state of techni-fermion and anti-techni-fermion, will be identified with the lowest KK mode

coming from the bulk scalar field Φ , not Φ_X . Thanks to the additional explicit bulk scalar field Φ_X , we naturally improve the matching with the OPE of the underlying theory (QCD and SWC-TC) for current correlators so as to reproduce gluonic $1/Q^4$ term, which is clearly distinguished from the same $1/Q^4$ terms from chiral condensate in the case of SWC-TC with $\gamma_m \simeq 1$. Our model with $\gamma_m = 0$ and $N_f = 3$ well reproduces the real-life QCD (See Table I). We find that the QCD ρ meson mass M_ρ includes a (negative) contribution about 10% from the gluon condensate.

We analyze a generic case with $0 \lesssim \gamma_m \lesssim 1$ and calculate masses of the techni- ρ meson (M_ρ), the techni- a_1 meson (M_{a_1}), and the flavor-singlet scalar meson, techni-dilaton (M_{TD}), as well as the S parameter. We discuss the general tendency of the dependence of the meson masses relative to F_π , (M_ρ/F_π , M_{a_1}/F_π , M_{TD}/F_π) on γ_m , S , and Γ . We find a characteristic feature of the techni-dilaton mass related to the conformality of SWC-TC: For fixed S and γ_m , (M_ρ/F_π) and (M_{a_1}/F_π) are not sensitive to Γ , while (M_{TD}/F_π) substantially decreases as Γ increases. Actually, in the formal limit $\Gamma \rightarrow \infty$, we would have $M_{\text{TD}}/F_\pi \rightarrow 0$ (This does not imply the existence of the isolated true massless NG boson of the scale symmetry, since in our case the decay constant F_{TD} diverges and the techni-dilaton gets decoupled in that limit, see text.) For fixed S and Γ , again (M_ρ/F_π) and (M_{a_1}/F_π) are not sensitive to γ_m , while (M_{TD}/F_π) substantially decreases as γ_m increases.

Particularly for the case of $\gamma_m = 1$, we study the dependence of the S parameter on (M_ρ/F_π) for typical values of Γ . It is shown that the techni-gluon contribution reduces the value of S maximally about 10% in the region of $\hat{S} \lesssim 0.1$, although the general tendency is similar to the previous paper [39] without techni-gluon condensation: $\hat{S}$ decreases monotonically with respect to (F_π/M_ρ) to continuously approach zero. This implies (M_ρ/F_π) necessarily increases when $\hat{S}$ is required to be smaller.

To be more concrete, we consider a couple of typical models of SWC-TC with $\gamma_m \simeq 1$ and $N_{\text{TC}} = 2, 3, 4$ based on the CBZ-IRFP in the large N_f QCD. Using some specific dynamical features of the conformal anomaly indicated by the analysis based on the ladder SD equation (Eqs.(4.3) and (4.4)), we find the relation of Γ to ($\Lambda_{\text{ETC}}/F_\pi$): In the case of $N_{\text{TC}} = 3$ ($N_f = 4N_{\text{TC}}$) and $S \simeq 0.1$, we have $\Gamma \simeq 7$ for ($\Lambda_{\text{ETC}}/F_\pi$) = 10^4 – 10^5 (required by the FCNC constraint). Thanks to the large anomalous dimension γ_m and large techni-gluon condensation Γ , we obtain a relatively light techni-dilaton with mass $M_{\text{TD}} \simeq 600$ GeV compared with $M_\rho \simeq M_{a_1} \simeq 3.8$ TeV, consistently with the perturbative unitarity of $W_L W_L$ scattering. Note that M_ρ and M_{a_1} are essentially determined by the requirement of $S = 0.1$ fairly independently of techni-gluon condensation. The essential reason for the large Γ is due to the existence of the wide conformal region $F_\pi < \mu < \Lambda_{\text{ETC}}$ with ($\Lambda_{\text{ETC}}/F_\pi$) = 10^4 – 10^5 , which yields the smallness of the beta function through the factor $(\ln 4\Lambda_{\text{ETC}}/\mu)^{-3}$ (see Eq.(4.4)) and hence amplifies the techni-gluon condensation compared with the ordinary QCD with $\Gamma = 1$. In the idealized (phenomenologically non-interesting) limit $\Lambda_{\text{ETC}}/F_\pi \rightarrow \infty$ we would have $\Gamma \rightarrow \infty$ and hence $M_{\text{TD}}/F_\pi \rightarrow 0$ in conformity with the general tendency mentioned above. (However, the would-be “massless” techni-dilaton is actually decoupled, see text. Indeed, spontaneous breaking of the scale symmetry is always accompanied with its explicit breaking.) The predicted mass $\simeq 600$ GeV of the holographic techni-dilaton (“conformal Higgs”) is within reach of LHC discovery.

This paper is organized as follows: In Sec. II we present our model which is an extension of our previous holographic SWC-TC model based on hard-wall-type bottom-up approach [39] by including effects of (techni-) gluon condensation through the bulk flavor/chiral-singlet scalar field. Formulas for masses of mesons (techni- ρ , $-a_1$, and -dilaton) and current correlators including S parameter are given. We show that our model reproduces gluonic $1/Q^4$ terms in the OPE of vector/axial-vector current correlators. In Sec. III we estimate effects on meson masses and S parameter coming from the (techni-)gluon condensation Γ as a free parameter in a generic TC with $0 \lesssim \gamma_m \lesssim 1$ involving the case of QCD with $\gamma_m \simeq 0$. In Sec. IV, to specify the value of Γ relevant to the actual model-building of SWC-TC, we consider a matching of our holographic model with typical models of SWC-TC based on the CBZ-IRFP in the large N_f QCD.

II. A HOLOGRAPHIC TECHNICOLOR MODEL WITH TECHNI-GLUON CONDENSATION

In this section, we propose a holographic model dual to a generic class of technicolor (TC) with $0 \lesssim \gamma_m \lesssim 1$ including the degree of freedom of (techni-)gluon condensation, where γ_m denotes the anomalous dimension of (techni-)fermion chiral condensate $\langle \bar{T}T \rangle$.

Following a bottom-up approach of holographic-dual of QCD [30, 31] with $\gamma_m \simeq 0$ and that of SWC-TC [37–39] with $\gamma_m \simeq 1$, we consider a five-dimensional gauge theory having $SU(N_f)_L \times SU(N_f)_R$ gauge symmetry. We will not consider the extra $U(1)_A$ that involves the anomaly. The theory is defined on the five-dimensional anti-de-Sitter space (AdS_5) with L , the curvature radius of AdS_5 , described by the metric $ds^2 = g_{MN}dx^M dx^N = (L/z)^2 (\eta_{\mu\nu}dx^\mu dx^\nu - dz^2)$ with $\eta_{\mu\nu} = \text{diag}[1, -1, -1, -1]$. The fifth direction z is compactified on an interval extended from the ultraviolet (UV) brane located at $z = \epsilon$ to the infrared (IR) brane at $z = z_m$, i.e., $\epsilon \leq z \leq z_m$. In addition to the bulk left- (L_M) and right- (R_M) gauge fields, we introduce a bulk scalar field Φ which transforms as bifundamental representation under the $SU(N_f)_L \times SU(N_f)_R$ gauge symmetry so as to deduce the information concerning the chiral condensation-operator

$\bar{T}T$. The mass-parameter m_Φ is then related to γ_m as $m_\Phi^2 = -(3 - \gamma_m)(1 + \gamma_m)/L^2$, where $\gamma_m = 0$ corresponds to QCD and QCD-like TC and $\gamma_m \simeq 1$ is the case of SWC-TC. This is the same setup as in Refs. [37–39].

In order to incorporate effects from techni-gluon condensation, here we introduce an additional bulk scalar field Φ_X dual to techni-gluon condensate $\langle \alpha G_{\mu\nu}^2 \rangle$, where α is related to the TC gauge coupling g_{TC} by $\alpha = g_{\text{TC}}^2/(4\pi)$. Since $\langle \alpha G_{\mu\nu}^2 \rangle$ is singlet under the chiral $SU(N_f)_L \times SU(N_f)_R$ symmetry and $U(1)_V$ symmetry, the dual-bulk scalar field Φ_X has to be a real field. We take $\dim(\alpha G_{\mu\nu}^2) = 4$ and the corresponding bulk-mass parameter $m_{\Phi_X}^2 = 0$.

The form of interaction terms involving the new bulk field Φ_X still remains undetermined. In the present work, we shall adopt a “dilaton-like” coupling, such that all the fields couple to Φ_X in the exponential form like $e^{\Phi_X(z)}$ ^{#3}. (Φ_X is *not* identified with techni-dilaton in this paper.)

Thus the five-dimensional action employed in the present paper takes the form:

$$S_5 = \int d^4x \int_\epsilon^{z_m} dz \sqrt{-g} \frac{1}{g_5^2} e^{c g_5^2 \Phi_X(z)} \left(-\frac{1}{4} \text{Tr} [L_{MN} L^{MN} + R_{MN} R^{MN}] \right. \\ \left. + \text{Tr} [D_M \Phi^\dagger D^M \Phi - m_\Phi^2 \Phi^\dagger \Phi] + \frac{1}{2} \partial_M \Phi_X \partial^M \Phi_X \right), \quad (2.1)$$

where $L_M(R_M) = L_M^a(R_M^a)T^a$ with the generators of $SU(N_f)$ normalized by $\text{Tr}[T^a T^b] = \delta^{ab}$; $L(R)_{MN} = \partial_M L(R)_N - \partial_N L(R)_M - i[L(R)_M, L(R)_N]$; $g = \det[g_{MN}] = -(L/z)^{10}$; g_5 denotes the gauge coupling in five-dimension and c is the dimensionless coupling constant. The covariant derivative acting on Φ is defined as $D_M \Phi = \partial_M \Phi + i L_M \Phi - i \Phi R_M$.

We parametrize the bulk scalar fields Φ and Φ_X as follows:

$$\Phi(x, z) = \frac{1}{\sqrt{2}}(v(z) + \sigma(x, z)) \exp[i\pi(x, z)/v(z)], \quad (2.2)$$

$$\Phi_X(z) = v_X(z), \quad (2.3)$$

with the vacuum expectation values (VEVs), $v(z) = \sqrt{2}\langle \Phi \rangle$ and $v_X(z) = \langle \Phi_X \rangle$, respectively. In Eq.(2.3) we ignored Kaluza-Klein (KK) modes of Φ_X (including the lowest mode) which are identified with massive glueballs with mass of order $\mathcal{O}(\Lambda_{\text{TC}})$ which is much larger than the electroweak scale, $\Lambda_{\text{TC}} \gg F_\pi$, in the case of SWC-TC with $\gamma_m \simeq 1$. The techni-dilaton, a flavor-singlet scalar bound state of techni-fermion and anti-techni-fermion, will be identified with the lowest KK mode of $\sigma(x, z)$, but not of Φ_X .

We choose the boundary condition for $v(z)$ as

$$\alpha M = \lim_{\epsilon \rightarrow 0} Z_m \left(\frac{L}{z} v(z) \right) \Big|_{z=\epsilon}, \quad Z_m = Z_m(L/z) = \left(\frac{L}{z} \right)^{\gamma_m}, \quad (2.4)$$

$$\xi = L v(z) \Big|_{z=z_m}, \quad (2.5)$$

where M stands for the current mass of techni-fermions and ξ is related to the techni-fermion condensate $\langle \bar{T}T \rangle$ as will be clarified later (See Eq.(2.27)). The parameter α has been introduced which can arise from ambiguity of definition of the current mass M . Here we take $\alpha = \sqrt{3}$, which turns out to be consistent with the operator product expansion (OPE) for the scalar current correlator in QCD [41].

For v_X , we impose the following boundary condition:

$$M' = \lim_{\epsilon \rightarrow 0} L v_X(z) \Big|_{z=\epsilon}, \quad \mathcal{G} = L v_X(z) \Big|_{z=z_m}, \quad (2.6)$$

where M' becomes the external source for the techni-gluon condensation-operator $\alpha G_{\mu\nu}^2$ and $\mathcal{G}$ is associated to the techni-gluon condensate $\langle \alpha G_{\mu\nu}^2 \rangle$ as we will see later (See Eq.(2.26)). We define the techni-gluon condensate in such a way that it does not include a trivial perturbative contribution, namely,

$$\langle \alpha G_{\mu\nu}^2 \rangle \equiv \langle \alpha G_{\mu\nu}^2 \rangle_{\text{full}} - \langle \alpha G_{\mu\nu}^2 \rangle_{\text{perturbation}}. \quad (2.7)$$

We then see that $\mathcal{G}$ in Eq.(2.6) is related only to the non-perturbative breaking of the conformal/scale invariance, while M' in Eq.(2.6) serves as its source and itself plays a role of the explicit breaking of the conformal symmetry just like the current mass M in the case of the chiral symmetry.

^{#3} This should not be confused with so-called soft-wall model [40] where $1/z_m = 0$ in contrast to our case with $1/z_m \neq 0$.

We next introduce the five-dimensional vector and axial-vector gauge fields V_M and A_M defined by $V_M = (L_M + R_M)/\sqrt{2}$ and $A_M = (L_M - R_M)/\sqrt{2}$. The UV boundary values of V_μ and A_μ then play the role of the sources (v_μ , a_μ) for the vector and the axial-vector currents externally coupled to TC sector. Working in $V_z = A_z \equiv 0$ gauge, we choose their boundary conditions as

$$\begin{aligned} \partial_z V_\mu(x, z)|_{z=z_m} &= \partial_z A_\mu(x, z)|_{z=z_m} = 0, \\ V_\mu(x, z)|_{z=\epsilon} &= v_\mu(x), \quad A_\mu(x, z)|_{z=\epsilon} = a_\mu(x). \end{aligned} \quad (2.8)$$

Once the boundary conditions (2.4), (2.5), (2.6), and (2.8) are given, the equations of motion for the bulk fields can be solved at the classical level. By substituting those solutions into the action (2.1), the effective action S_5^{eff} is expressed as a certain functional of the UV boundary values, M , M' , v_μ , and a_μ , i.e., $S_5^{\text{eff}} = S_5^{\text{eff}}[M, M', v_\mu, a_\mu]$. From the familiar AdS/CFT dictionary, this S_5^{eff} corresponds to the generating functional W in TC written in terms of the external sources M , M' , v_μ , and a_μ . One can then readily calculate the two-point Green functions in the usual way:

$$\left. \frac{\delta^2 W[v_\mu]}{\delta \tilde{v}_\mu^a(q) \delta \tilde{v}_\nu^b(-q)} \right|_{v_\mu=0} = i \int d^4x e^{iq \cdot x} \langle J_V^{a\mu}(x) J_V^{b\nu}(0) \rangle = -\delta^{ab} \left(\eta^{\mu\nu} - \frac{q^\mu q^\nu}{q^2} \right) \Pi_V(-q^2), \quad (2.9)$$

$$\left. \frac{\delta^2 W[a_\mu]}{\delta \tilde{a}_\mu^a(q) \delta \tilde{a}_\nu^b(-q)} \right|_{a_\mu=0} = i \int d^4x e^{iq \cdot x} \langle J_A^{a\mu}(x) J_A^{b\nu}(0) \rangle = -\delta^{ab} \left(\eta^{\mu\nu} - \frac{q^\mu q^\nu}{q^2} \right) \Pi_A(-q^2), \quad (2.10)$$

$$\lim_{\epsilon \rightarrow 0} i \frac{\delta W[M]}{\delta M} \Big|_{M=0} = \langle \bar{T}T \rangle, \quad (2.11)$$

$$\lim_{\epsilon \rightarrow 0} i \frac{\delta W[M']}{\delta M'} \Big|_{M'=0} = \langle \alpha G_{\mu\nu}^2 \rangle, \quad (2.12)$$

where $\tilde{v}^\mu(q)$ and $\tilde{a}^\mu(q)$ respectively denote the Fourier component of $v^\mu(x)$ and $a^\mu(x)$. To facilitate the later discussions, as done in Ref. [39], we introduce $\hat{S}$ as the S parameter per each techni-fermion doublet, $\hat{S} = S/(N_f/2)$, which is expressed by the vector and axial-vector current correlators Π_V and Π_A as

$$\hat{S} = -4\pi \frac{d}{dQ^2} [\Pi_V(Q^2) - \Pi_A(Q^2)]_{Q^2=0}, \quad (2.13)$$

where $Q \equiv \sqrt{-q^2}$. We also introduce the (techni-)pion decay constant defined as

$$F_\pi^2 = \Pi_V(0) - \Pi_A(0). \quad (2.14)$$

A. Condensates: $\langle \bar{T}T \rangle$ and $\langle \alpha G_{\mu\nu}^2 \rangle$

In the following, evaluating the equations of motion for the bulk fields explicitly, we shall present the formulas for the condensates, $\langle \bar{T}T \rangle$ and $\langle \alpha G_{\mu\nu}^2 \rangle$, the S parameter and the decay constant F_π , and masses of vector mesons, axial-vector mesons, and scalar mesons.

The action (2.1) leads to the following equation of motion for $v_X(z)$:

$$\partial_z \left(\frac{1}{z^3} \partial_z \chi(z) \right) + \frac{(cg_5^2)^2}{4L^2} \chi(z) L^2 \text{Tr} \left[-\frac{1}{z^3} (\partial_z v(z))^2 + \frac{(3 - \gamma_m)(1 + \gamma_m)}{z^5} v^2(z) \right] = 0, \quad (2.15)$$

where we have defined

$$\chi(z) = \exp \left[\frac{c}{2} g_5^2 v_X(z) \right]. \quad (2.16)$$

The boundary condition for Φ_X given in Eq.(2.6) is now rewritten in terms of χ as

$$\chi(z) \Big|_{z=\epsilon} = \exp \left[\frac{c}{2} g_5^2 v_X(\epsilon) \right] = \exp \left[\frac{c}{2} \frac{g_5^2}{L} M' \right], \quad (2.17)$$

$$\chi(z) \Big|_{z=z_m} = \exp \left[\frac{c}{2} g_5^2 v_X(z_m) \right] = \exp \left[\frac{c}{2} \frac{g_5^2}{L} \mathcal{G} \right] \equiv G + 1. \quad (2.18)$$

We solve the equation of motion (2.15) keeping only the first term of the left hand side in Eq.(2.15). This assumption will be justified later in determining the size of $|(c/2)(g_5^2/L)|^2$, which turns out to be $\simeq 10^{-4}$ (See Eq.(3.2)). Equation (2.15) is now easily solved to give the solution

$$\chi(z) = \exp \left[\frac{c}{2} g_5^2 v_X(z) \right] = c_1^\chi + c_2^\chi \left(\frac{z}{L} \right)^4, \quad (2.19)$$

where c_1^χ and c_2^χ are determined by Eqs.(2.17) and (2.18) in the limit $\epsilon \rightarrow 0$ as

$$c_1^\chi = e^{\frac{c}{2} \frac{g_5^2}{L} M'}, \quad c_2^\chi = \left(\frac{L}{z_m} \right)^4 (G + 1 - c_1^\chi). \quad (2.20)$$

Note that the solution in Eq.(2.19) gives rise to the induced metric for the vector and axial-vector gauge fields (See Eq.(2.29)).

We next turn to the equation of motion for $v(z)$ which is read off from the action (2.1) as follows:

$$\partial_z \left(\frac{1}{z^3} \chi^2(z) \partial_z v(z) \right) + \chi^2(z) \frac{(3 - \gamma_m)(1 + \gamma_m)}{z^5} v(z) = 0. \quad (2.21)$$

Substituting Eq.(2.19) into Eq.(2.21) and taking $M' = 0$, we find the solution in the limit $\epsilon \rightarrow 0$

$$v(z) = \frac{1}{1 + G \left(\frac{z}{z_m} \right)^4} \left(c_1 \left(\frac{z}{L} \right)^{\gamma_m + 1} + c_2 \left(\frac{z}{L} \right)^{3 - \gamma_m} \right), \quad (2.22)$$

where c_1 and c_2 are determined by the boundary condition in Eqs.(2.4) and (2.5) as

$$c_1 = \sqrt{3} M, \quad (2.23)$$

$$c_2 = \frac{\xi(1 + G)}{L} \left(\frac{L}{z_m} \right)^{3 - \gamma_m} - \left(\frac{L}{z_m} \right)^{2(1 - \gamma_m)} c_1. \quad (2.24)$$

Note that in Eqs.(2.22) and (2.24) (techni-)gluon condensation effects are included: When $G = 0$ in Eqs.(2.22) and (2.24), we get back to the previous results without (techni-)gluon-condensation effects [30, 31, 37–39].

Putting the classical solutions, Eqs.(2.19) and (2.22), into the action (2.1), we are left with the four-dimensional boundary term which is holographically dual to the generating functional $W[M, M']$ in TC,

$$W[M, M'] = \int d^4x \frac{L^3}{2g_5^2} \left[-\frac{1}{z^3} \frac{4}{(cg_5^2)^2} \partial_z \chi(z) \cdot \chi(z) - \frac{1}{z^3} \chi^2(z) \text{Tr} [\partial_z v(z) \cdot v(z)] \right]_\epsilon^{z_m}. \quad (2.25)$$

Using Eqs.(2.11) and (2.12) and performing the functional derivative with respect to the sources M and M' , the techni-gluon condensate $\langle \alpha G_{\mu\nu}^2 \rangle$ and the techni-fermion condensate $\langle \bar{T}T \rangle$ are respectively expressed in terms of the five-dimensional gauge theory as

$$\langle \alpha G_{\mu\nu}^2 \rangle = -8 \frac{L^2}{cg_5^4} \frac{1}{z_m^4} G, \quad (2.26)$$

$$\langle \bar{T}T \rangle_{1/L} = -\sqrt{3} \frac{L}{g_5^2} \frac{(3 - \gamma_m)}{z_m^3} (1 + G) \xi \cdot Z_m^{-1}, \quad (2.27)$$

where $Z_m = (L/z_m)^{\gamma_m}$.

B. Vector, axial-vector current correlators, decay constant F_π , and S parameter

Let us next focus on the vector and axial-vector sectors. The relevant action under the gauge-fixing $V_z = A_z \equiv 0$ reads

$$S_5 \ni -\frac{1}{2g_5^2} \int d^4x \int_\epsilon^{z_m} dz w(z) \left(\text{Tr} \left[\frac{1}{2} V_{\mu\nu} V^{\mu\nu} - \partial_z V_\mu \partial_z V^\mu + \frac{1}{2} A_{\mu\nu} A^{\mu\nu} - \partial_z A_\mu \partial_z A^\mu \right] - 2 \left(\frac{L}{z} \right)^2 \text{Tr} [v^2(z) A_\mu A^\mu] \right), \quad (2.28)$$

where $V(A)_{\mu\nu} = \partial_\mu V(A)_\nu - \partial_\nu V(A)_\mu$ and the induced metric $w(z)$ is given by the solution in Eq.(2.19) as

$$w(z) = \frac{L}{z} \chi^2(z) = \frac{L}{z} \left(1 + G \left(\frac{z}{z_m} \right)^4 \right)^2. \quad (2.29)$$

In arriving at the last equality we have used Eqs.(2.19), (2.20), and set $M' = 0$. (When $M' = 0$ and $\epsilon = 0$ the explicit breaking of conformal/scale invariance only comes from $1/z_m \neq 0$.) The action in Eq.(2.28) takes the same form as in Refs. [30, 31, 37–39] except that $w(z) = L/z$ has been replaced by the one given in Eq.(2.29). Our induced metric determined by the equation of motion for the bulk scalar Φ_X is compared with the effective metric of Ref. [42] lacking the bulk scalars Φ and Φ_X , where the form of the effective metric was simply assumed to reproduce the OPE for the vector and axial-vector current correlators $\Pi_{V,A}$.

We solve the equations of motion for the transversely polarized components of the gauge fields $V_\mu(x, z)$ and $A_\mu(x, z)$. The corresponding equations of motion are immediately read off from the action (2.28) as

$$[q^2 + w(z)^{-1} \partial_z w(z) \partial_z] V_\mu(q, z) = 0, \quad (2.30)$$

$$\left[q^2 + w(z)^{-1} \partial_z w(z) \partial_z - 2 \left(\frac{L}{z} \right)^2 v^2(z) \right] A_\mu(q, z) = 0, \quad (2.31)$$

where $V_\mu(q, z)$ and $A_\mu(q, z)$ denote the Fourier transforms of $V_\mu(x, z)$ and $A_\mu(x, z)$, respectively. It is convenient to decompose $V_\mu(q, z)$ and $A_\mu(q, z)$ into the external sources $(\tilde{v}_\mu(q), \tilde{a}_\mu(q))$ and the remainders $(V(q, z), A(q, z))$, such as $V_\mu(q, z) = \tilde{v}_\mu(q) V(q, z)$ and $A_\mu(q, z) = \tilde{a}_\mu(q) A(q, z)$. Using the equations of motion (2.30) and (2.31) together with the boundary conditions in Eq.(2.8), we rewrite the action (2.28) to get the four-dimensional UV boundary term which is holographically dual to the generating functional $W[v_\mu, a_\mu]$ in TC,

$$W[v_\mu, a_\mu] = \frac{1}{2} \int \frac{d^4 q}{(2\pi)^4} \frac{-1}{g_5^2} w(\epsilon) \text{Tr} [\tilde{v}_\mu(-q) \partial_z V(q, \epsilon) \cdot \tilde{v}^\mu(q) + \tilde{a}_\mu(-q) \partial_z A(q, \epsilon) \cdot \tilde{a}^\mu(q)], \quad (2.32)$$

where $V(q, z)$ and $A(q, z)$ respectively satisfy the boundary conditions,

$$V(q, \epsilon) = 1, \quad \partial_z V(q, z_m) = 0, \quad (2.33)$$

$$A(q, \epsilon) = 1, \quad \partial_z A(q, z_m) = 0. \quad (2.34)$$

From Eqs.(2.9) and (2.10), the vector and axial-vector current correlators Π_V and Π_A are now expressed in terms of the five-dimensional gauge theory as

$$\Pi_V(Q^2) = \frac{w(\epsilon)}{g_5^2} \partial_z V(Q^2, \epsilon), \quad \Pi_A(Q^2) = \frac{w(\epsilon)}{g_5^2} \partial_z A(Q^2, \epsilon), \quad (2.35)$$

where we have rewritten $V(q, z) = V(Q^2, z)$ and $A(q, z) = A(Q^2, z)$. We emphasize that, thanks to the introduction of the bulk scalar field Φ_X dual to the role of the gluon condensation, the present model reproduces $\langle \alpha G_{\mu\nu}^2 \rangle / Q^4$ term in the OPE of the current correlators Π_V and Π_A , which was missing in the previous approach without Φ_X [30, 31, 37–39]. Leaving the details of the derivation in Appendix A, we will here just show the resultant expression of the high-energy expansion of $\Pi_{V,A}(Q^2)$ in the large Euclidean-momentum region $(1/z_m)^2 \ll Q^2 < (1/\epsilon)^2$,

$$\Pi_{V,A}(Q^2) \Big|_{(1/z_m)^2 \ll Q^2 < (1/\epsilon)^2} = Q^2 \left[\frac{L}{2g_5^2} \ln Q^2 + c \frac{2}{3} \frac{g_5^2}{L} \frac{\langle \alpha G_{\mu\nu}^2 \rangle}{Q^4} + \mathcal{O}\left(\frac{1}{Q^{6-2\gamma_m}}\right) \right]. \quad (2.36)$$

Furthermore, by introducing some extra higher-dimensional interaction terms (See Eq.(A.8)), the present model exactly reproduces the high-energy behavior up to terms suppressed by $(1/Q^8)$, consistently with the form of the OPE,

$$\Pi_{V,A}(Q^2) \Big|_{(1/z_m)^2 \ll Q^2 < (1/\epsilon)^2} = Q^2 \left[\frac{L}{2g_5^2} \ln Q^2 + c \frac{2}{3} \frac{g_5^2}{L} \frac{\langle \alpha G_{\mu\nu}^2 \rangle}{Q^4} + C_6^{V,A} L^{2\gamma_m} \frac{\langle \bar{T} T \rangle_{1/L}^2}{Q^{2(3-\gamma_m)}} + \mathcal{O}\left(\frac{1}{Q^8}\right) \right], \quad (2.37)$$

where the couplings $C_6^{V,A}$ come from the higher-dimensional interaction terms (See Eq.(A.14)). It should also be stressed that such extra interaction terms do not affect all of our results shown in the later sections.

Our model is sharply contrasted with the approach in Ref. [42] where the effective metric was assumed so as to produce $1/Q^4$ term which, however, could be confused with the chiral condensation term $\langle \bar{T}T \rangle^2/Q^{6-2\gamma_m} \simeq \langle \bar{T}T \rangle^2/Q^4$ in the case of SWC-TC with $\gamma_m \simeq 1$ in the absence of the bulk scalars Φ and Φ_X . In our case which explicitly includes the bulk scalar field Φ_X dual to the gluon condensate, we are able to obtain not just the form behaving as $(1/Q^4)$ but the whole expression $\langle \alpha G_{\mu\nu}^2 \rangle / Q^4$ involving the gluon condensate $\langle \alpha G_{\mu\nu}^2 \rangle$, and hence clearly distinguish from the $\langle \bar{T}T \rangle^2/Q^4$ term arising due to the bulk scalar Φ .

In order to obtain the formulas for the decay constant F_π and the S parameter, we shall expand $\Pi_{V,A}(Q^2)$ perturbatively in powers of Q^2 as $\Pi_{V,A}(Q^2) = \Pi_{V,A}(0) + Q^2 \Pi'_{V,A}(0) + \mathcal{O}(Q^4)$, where $\Pi'_{V,A}(0) \equiv \partial \Pi_{V,A}(Q^2) / \partial Q^2 \Big|_{Q^2=0}$. Then $\Pi_{V,A}(0)$ and $\Pi'_{V,A}(0)$ are expressed as

$$\Pi_V(0) = 0, \quad \Pi'_V(0) = -\frac{L}{g_5^2} \int_\epsilon^{z_m} \frac{dz'}{z'} \chi^2(z'), \quad (2.38)$$

$$\Pi_A(0) = \frac{L}{g_5^2} \frac{1}{\epsilon} \chi^2(\epsilon) \partial_z A(0, z) \Big|_{z=\epsilon}, \quad \Pi'_A(0) = -\frac{L}{g_5^2} \int_\epsilon^{z_m} \frac{dz'}{z'} \chi^2(z') A^2(0, z'), \quad (2.39)$$

where $A(0, z)$ is given as a solution to Eq.(2.31) with $q^2 = 0$. From Eqs.(2.14) and (2.13), we find that F_π and $\hat{S}$ are expressed in terms of the five-dimensional gauge theory as

$$F_\pi^2 = -\frac{L}{g_5^2} \frac{1}{\epsilon} \chi^2(\epsilon) \partial_z A(0, z) \Big|_{z=\epsilon}, \quad (2.40)$$

$$\hat{S} = 4\pi \frac{L}{g_5^2} \int_\epsilon^{z_m} \frac{dz}{z} \chi^2(z) (1 - A^2(0, z)). \quad (2.41)$$

C. Vector, axial-vector, and flavor-singlet scalar meson masses: M_{V_n} , M_{A_n} , and M_{σ_n}

A set of vector meson masses $\{M_{V_n}\}$ arises as an infinite tower of eigenvalues of normalizable solutions $\{V_n(z)\}$ satisfying Eq.(2.30) with q^2 replaced by $\{M_{V_n}^2\}$,

$$\left[M_{V_n}^2 + w(z)^{-1} \partial_z w(z) \partial_z \right] V_n(z) = 0, \quad (2.42)$$

with the boundary condition $V_n(\epsilon) = 0$ and $\partial_z V_n(z_m) = 0$. The lowest eigenvalue is identified as the techni- ρ meson mass, $M_{V_1} = M_\rho$.

Similarly for axial-vector meson masses $\{M_{A_n}\}$, the eigenvalue equation for a set of normalizable modes $\{A_n(z)\}$ is obtained by taking $q^2 = M_{A_n}^2$ in Eq.(2.31):

$$\left[M_{A_n}^2 + w(z)^{-1} \partial_z w(z) \partial_z - 2 \left(\frac{L}{z} \right)^2 v^2(z) \right] A_n(z) = 0, \quad (2.43)$$

with the boundary condition $A_n(\epsilon) = 0$ and $\partial_z A_n(z_m) = 0$. The lowest eigenvalue is regarded as the techni- a_1 meson mass, $M_{A_1} = M_{a_1}$.

The equation of motion for the flavor-singlet scalar field $\sigma(x, z)$ is decomposed into the eigenvalue equations for a set of the KK modes $\sigma^{(n)}(x)$ arising as $\sigma(x, z) = \sum_{n=1}^\infty \sigma^{(n)}(x) \sigma_n(z)$. By taking into account Eq.(2.2) and replacing the momentum-squared q^2 with the mass-squared $M_{\sigma_n}^2$, the equation of motion for the wave function $\sigma_n(z)$ is read off from the action (2.1) as

$$\left[M_{\sigma_n}^2 + \left(\frac{w(z)}{z^2} \right)^{-1} \partial_z \left(\frac{w(z)}{z^2} \right) \partial_z - \frac{(3 - \gamma_m)(1 + \gamma_m)}{z^2} \right] \sigma_n(z) = 0, \quad (2.44)$$

where the normalizable solution $\sigma_n(z)$ should satisfy the UV boundary condition, $\lim_{\epsilon \rightarrow 0} \sigma_n(\epsilon) = 0$, so as to make the action finite at $z = \epsilon$ in the limit $\epsilon \rightarrow 0$. The solution to Eq.(2.44) is then given as

$$\sigma_n(z) = c_\sigma \frac{z^2}{1 + G\left(\frac{z}{z_m}\right)} J_{1-\gamma_m}(M_{\sigma_n} z), \quad (2.45)$$

which satisfies the UV boundary condition. In Eq.(2.45) J_n is Bessel function of order n and the overall coefficient c_σ is determined by a normalization condition which is irrelevant to the present study.

The eigenvalue equation for M_{σ_n} is provided by the IR boundary condition which needs a bit careful consideration because it is related to a vacuum stabilization problem: The vacuum energy on the background of the bulk scalar field Φ is equivalent to the generating functional in Eq.(2.25). By using the solution for $v(z)$ in Eq.(2.22), the vacuum energy is expressed as a function of ξ in the chiral limit $M = 0$,

$$V[\xi] = - \int d^4x \frac{L}{2g_5^2} \chi^2(z_m) \text{Tr} \left[\left(3 - \gamma_m - 4 \frac{G}{1+G} \right) \frac{\xi^2}{z_m^4} \right], \quad (2.46)$$

where we have put $M' = 0$ for simplicity. One can easily see that, with respect to ξ , $V[\xi]$ is minimized at $\xi = 0$, which readily leads to $\langle \bar{T}T \rangle = 0$ through Eq.(2.27) and hence to no spontaneous breaking of chiral symmetry. In order to avoid this problem, similarly to a procedure proposed in Ref. [41], we may introduce the following IR potential:

$$\begin{aligned} \mathcal{L}_{\text{IR}} &= - \left(\frac{L}{z} \right)^4 \chi^2(z) V(\Phi) \Big|_{z_m}, \\ V(\Phi)|_{z=z_m} &= -m_b^2 \text{Tr} |\Phi(z_m)|^2 + \lambda \text{Tr} (|\Phi(z_m)|^2)^2, \end{aligned} \quad (2.47)$$

where the potential parameters m_b^2 and λ are taken to be positive. By adding this IR potential, the vacuum is now realized at $\xi \neq 0$:

$$\xi^2 = \frac{1}{\lambda} \left(L^2 m_b^2 - \frac{L}{g_5^2} \left(3 - \gamma_m - 4 \frac{G}{1+G} \right) \right), \quad (2.48)$$

where ξ^2 is tuned to be positive by adjusting m_b^2 . The IR boundary condition for $\sigma_n(z)$ is now assigned in a way that the total IR boundary term with respect to $\sigma^{(n)}(x)$ is canceled in the quadratic order:

$$\left[\partial_z + 2 \left(\frac{L}{z_m} \right) g_5^2 m_{\sigma_5}^2 \right] \sigma_n(z) \Big|_{z_m} = 0, \quad (2.49)$$

where

$$m_{\sigma_5}^2 = \frac{1}{L^2} \left[\lambda \xi^2 - \frac{1}{2} \frac{L}{g_5^2} \left(3 - \gamma_m - 4 \frac{G}{1+G} \right) \right]. \quad (2.50)$$

Substituting the solution in Eq.(2.45) into the IR boundary condition (2.49), we thus obtain the eigenvalue equation for M_{σ_n} ,

$$2\lambda \xi^2 \frac{g_5^2}{L} J_{1-\gamma_m}(M_{\sigma_n} z_m) = M_{\sigma_n} z_m \cdot J_{2-\gamma_m}(M_{\sigma_n} z_m). \quad (2.51)$$

The lowest eigenvalue is identified as the techni-dilaton mass, $M_{\sigma_1} = M_{\text{TD}}$.

III. ANALYSIS ON GLUONIC-CONTRIBUTIONS

In this section, we shall discuss effects on observables coming from the gluon condensation in a generic TC with $0 \lesssim \gamma_m \lesssim 1$ involving the case of QCD with $\gamma_m \simeq 0$. Among observables, we particularly focus on the S parameter and the masses of the lowest KK modes for the vector, axial-vector, and flavor-singlet scalar mesons ($M_\rho, M_{a_1}, M_{\text{TD}}$). To this end, we first pay our attention to the parameters describing the present five-dimensional model and momentarily discuss how they can be fixed by considering the holographic-dual of the generic TC with $0 \lesssim \gamma_m \lesssim 1$. The parameters are following ten:

$$\frac{L}{g_5^2}, \quad z_m, \quad \epsilon, \quad \gamma_m, \quad \xi, \quad M, \quad G(\text{or } \mathcal{G}), \quad M', \quad c, \quad \lambda. \quad (3.1)$$

The UV brane position ϵ is treated as the cutoff scale ($1/\epsilon$) of the five-dimensional theory and is usually set to be 0 after all calculations are done. From a point of view of a typical TC scenario, on the other hand, the UV cutoff ($1/\epsilon$) can be replaced by an ETC scale ($1/\epsilon = \Lambda_{\text{ETC}}$). As for the IR brane position z_m , similarly, it can

play a role of the IR cutoff scale of the theory associated with the chiral symmetry breaking or confinement, and hence $(1/z_m)$ can be related to a typical meson mass scale, say, m , in TC. Since $(m/\Lambda_{\text{ETC}}) \ll 1$, we may simply put $(\epsilon/z_m) = (m/\Lambda_{\text{ETC}}) = 0$. Then we see that the S parameter does not depend on either ϵ or z_m since it is a dimensionless quantity. Note, however, that other dimensionful quantities, such as F_π , M_ρ , M_{a_1} , and M_{TD} , still have a certain z_m -dependence which can be completely factorized by defining dimensionless ones like $\tilde{F}_\pi = z_m F_\pi$, and so on.

The parameter M' is the external source of the techni-gluon condensation-operator $\alpha G_{\mu\nu}^2$ and hence is regarded as the explicit breaking source of the conformal/scale symmetry associated with the dilatation current anomaly characterized by the intrinsic scale of order Λ_{TC} ($\simeq \Lambda_{\text{ETC}} \gg 1/z_m$ for $\gamma_m \simeq 1$) in which we are not interested. Here we take $M' = 0$. (Even when $M' = 0$ and $\epsilon = 0$, we have the explicit breaking of conformal/scale invariance due to $1/z_m \neq 0$.)

The parameters (L/g_5^2) and c are determined by comparing the high-energy behavior of the current correlators $\Pi_{V,A}$ to those obtained by the OPE: (L/g_5^2) from the $\ln Q^2$ term and c from the $\langle \alpha G_{\mu\nu}^2 \rangle / Q^4$ term. In the case of SWC-TC with $\gamma_m \simeq 1$, $\langle \bar{T}T \rangle^2 / Q^4$ term has the same Q^2 -dependence as that of $\langle \alpha G_{\mu\nu}^2 \rangle / Q^4$ term. As was discussed in Sec. II B, it is possible to clearly distinguish those two terms in our approach. We will leave the detailed calculation in Appendix A and here just quote the result on the OPE-matching:

$$\frac{L}{g_5^2} = \frac{N_{\text{TC}}}{12\pi^2}, \quad c = -\frac{N_{\text{TC}}}{192\pi^3}. \quad (3.2)$$

The parameter λ has been introduced so as to minimize the bulk scalar potential with non-zero ξ . In order to know more about λ , let us take a look at the two terms in the square bracket of Eq.(2.50). One then finds that the first term ($\lambda\xi^2$) should be proportional to N_{TC} because of Eq.(3.2). Furthermore, from Eqs.(2.27) and (3.2) and taking into account $\langle \bar{T}T \rangle \propto N_{\text{TC}}$, we see that $\xi \propto N_{\text{TC}}^0$ and hence $\lambda \propto N_{\text{TC}}$. Supposing that the coupling λ is expected to be generated at one-loop level (through techni-fermion loops), we may totally write $\lambda = \kappa \times N_{\text{TC}} / (4\pi)^2$ with an $\mathcal{O}(1)$ parameter κ :

$$\lambda = \kappa \frac{N_{\text{TC}}}{(4\pi)^2}, \quad \kappa = 1.0 (\pm 0.3), \quad (3.3)$$

which reproduces the mass of the flavor-singlet scalar bound-state (two-quark state), $f_0(1370)$, in QCD as the lightest KK mode of the flavor-singlet scalar, M_{σ_1} . Actually in QCD, there are other two candidates for the light flavor-singlet scalar particles other than $f_0(1370)$, which are $f_0(600)$ (so-called σ) and $f_0(980)$. The following is the reason why we have adopted $f_0(1370)$ to fit the value of M_{σ_1} : Since we are interested in application to a generic TC, we need to carefully select a certain appropriate flavor-singlet scalar bound-state realized in a generic strongly coupled dynamics with arbitrary N_c and N_f . Thinking about the other two candidates from this point of view, one finds that those bound-states can be considered as four-quark states due to a specific property arising only in the case of real life QCD with $N_f = N_c = 3$ (See, for example, Ref. [43]), so they are excluded from candidate of two-quark state. We further notice that similar characteristic features in real life QCD would cause mixing between two-quark states and four-quark states, which could make the observed mass of the two-quark state $f_0(1370)$ lifted up. Without such a mixing, the mass of $f_0(1370)$ is expected to be around 1.1-1.2 GeV [44] ^{#4}. Thus we have determined the values of κ as in Eq.(3.3) ($0.7 \leq \kappa \leq 1.3$) using a set of QCD-fit values (See Eq(3.5)) so as to reproduce the allowed range of the mass of $f_0(1370)$ without the mixing.

Thus we are now left with the four undetermined parameters,

$$z_m, \quad \xi, \quad G, \quad \gamma_m (0 \lesssim \gamma_m \lesssim 1), \quad (3.4)$$

with z_m being the only dimensionful parameter.

A. QCD case with $\gamma_m \simeq 0$

We shall first consider the case of QCD with $N_c = 3$ and $\gamma_m \simeq 0$. In the case of QCD, the parameters ξ , G , and $(1/z_m)$ can be fixed in such a way that Eqs.(2.40), (2.42), and (2.26) respectively reproduce the experimental

^{#4} Actually in Ref.[44], the mixing between $a_0(980)$ and $a_0(1445)$ is discussed. It is known, however, that in the large N_c limit $f_0(1370)$ and $a_0(1445)$ are degenerate, so we can apply the similar argument against the mixing between $f_0(980)$ and $f_0(1370)$.

QCD-fit	M_{a_1}	$M_{\sigma_1=f_0(1370)}$	$\hat{S} = -16\pi L_{10}$	$(-\langle\bar{q}q\rangle)^{1/3}$
this model	1264 MeV	$1.15^{(+0.04)}_{(-0.07)} \text{ GeV}$	0.31	277 MeV
measured	$1230 \pm 40 \text{ MeV}$ [45]	*1.1-1.2 GeV	0.33 ± 0.04 [34]	$225 \pm 25 \text{ MeV}$ [47]

TABLE I: Values of several observables in QCD predicted from the present holographic model. The starred measured value of $M_{\sigma_1=f_0(1370)}$ corresponds to the mass estimated without mixing with a four quark state $f_0(980)$ (See discussion below Eq.(3.3)).

values [45] $f_\pi \simeq 92.4 \text{ MeV}$, $M_\rho \simeq 775 \text{ MeV}$, and a typical empirical value [46] $\frac{1}{\pi}\langle\alpha G_{\mu\nu}^2\rangle \simeq 0.012 \text{ GeV}^4$:

$$\xi \simeq 3.1, \quad G \simeq 0.25, \quad \frac{1}{z_m} \simeq 347 \text{ MeV}. \quad (3.5)$$

Using these values, we calculate the values of M_{a_1} (from Eq.(2.43)), $M_{\sigma_1} \equiv M_{f_0(1370)}$ (from Eq.(2.51) and using the values of κ in Eq.(3.3)), $\hat{S} (= -16\pi L_{10})$ (from Eq.(2.41)), and the quark condensate $\langle\bar{q}q\rangle$ (from Eq.(2.27)), which results in good agreement with experiment as shown in Table. I.

In our model, meson masses have some contributions from the gluon condensation. The effect of the gluon condensation on M_ρ can be analytically estimated for $G \simeq 0.25$ by expanding Eq.(2.42) perturbatively in G :

$$\begin{aligned} M_\rho &\simeq \frac{2.41}{z_m} \times [1 - 0.388 G] \\ &\simeq 836 \text{ MeV} \times [(1.000)_{G=0} - (0.097)_{G \neq 0}] \simeq 775 \text{ MeV}, \end{aligned} \quad (3.6)$$

where in reaching the last line we have used $(1/z_m) \simeq 347 \text{ MeV}$ in Eq.(3.5). Equation (3.6) implies that the gluon condensation negatively contributes about 10% to the ρ meson mass in QCD. Similar expression was obtained in somewhat different approach [42].

B. Generic TC case with $0 \lesssim \gamma_m \lesssim 1$

We next discuss the case of a generic TC with $0 \lesssim \gamma_m \lesssim 1$ involving SWC-TC with $\gamma_m \simeq 1$ ^{#5} and evaluate gluonic-contributions to the S parameter ($S = \hat{S} \cdot (N_f/2)$), masses of techni- ρ (M_ρ), techni- a_1 (M_{a_1}), and the lightest flavor-singlet scalar, techni-dilaton ($M_{\text{TD}} = M_{\sigma_1}$).

As we noted in Eq.(3.4), the dimensionless quantity $\hat{S}$ in Eq.(2.41) is given as a function of the three dimensionless parameters ξ , γ_m , and G ,

$$\hat{S} = \hat{S}(\gamma_m, \xi, G). \quad (3.7)$$

Other dimensionful quantities, F_π , M_ρ , M_{a_1} , and M_{TD} are respectively expressed as follows:

$$F_\pi = z_m^{-1} \cdot \widetilde{F}_\pi(\gamma_m, \xi, G), \quad (3.8)$$

$$M_\rho = z_m^{-1} \cdot \widetilde{M}_\rho(G), \quad (3.9)$$

$$M_{a_1} = z_m^{-1} \cdot \widetilde{M}_{a_1}(\gamma_m, \xi, G), \quad (3.10)$$

$$M_{\text{TD}} = z_m^{-1} \cdot \widetilde{M}_{\text{TD}}(\gamma_m, \xi; \kappa). \quad (3.11)$$

As in QCD, we will take the value of κ in Eq.(3.3), $\kappa = 1.0(\pm 0.3)$. Absence of explicit dependence of ξ and γ_m in M_ρ can be seen in Eq.(2.42). Explicit ξ - and γ_m -dependences for M_{a_1} enter in the $v^2(z)$ term in Eq.(2.43). M_{TD} has no explicit G -dependence as seen in Eq.(2.51). Note that every dimensionful quantities scales with the parameter $(1/z_m)$. Hereafter we shall consider dimensionless quantities (M_ρ/F_π) , (M_{a_1}/F_π) , and (M_{TD}/F_π) which are free from z_m .

^{#5} One might think that walking/near conformal dynamics is characterized by not $\gamma_m = 1$ but $\gamma_m \simeq 1$, as in a typical example of SWC-TC based on the Caswell-Banks-Zaks infrared fixed point [16, 17] in the large N_f QCD. However, as was clarified in Ref. [39], in the present holographic approach, there exists no discontinuity between $\gamma_m = 1$ and the limit $\gamma_m \rightarrow 1$, so that both cases give the same result. In the present work, therefore, we have explicitly taken $\gamma_m = 1$.




FIG. 2: Plots of (M_ρ/F_π) , (M_{a_1}/F_π) , and (M_{TD}/F_π) as a function of Γ with $N_{TC} = 3$ and $\hat{S} = 0.31$ fixed. The left panel and the right panel respectively correspond to the cases with $\gamma_m \simeq 0$ and $\gamma_m = 1$. In both panels, the dotted (red), dashed (yellow), and solid (blue) curves respectively denote (M_ρ/F_π) , (M_{a_1}/F_π) , and (M_{TD}/F_π) . In the left panel, the values on the vertical axis ($\Gamma = 1$) correspond to those of QCD.

To properly evaluate effects from the gluon condensate $\langle \alpha G_{\mu\nu}^2 \rangle$ in Eq.(2.26), we need to be a bit careful because the expression (2.26) involves not only G , the source of the gluon condensation, but also the IR cutoff scale $(1/z_m)$. This implies that a naive variation of the value of $\langle \alpha G_{\mu\nu}^2 \rangle$ would not merely lead to the change of the value of the gluon condensation itself. In order to extract the gluonic-contribution only, therefore, we shall use $\langle \alpha G_{\mu\nu}^2 \rangle / F_\pi^4$ which is not including $(1/z_m)$. It is convenient, furthermore, to work on the quantity normalized by the QCD value:

$$\Gamma = \Gamma(\gamma_m, \xi, G) \equiv \left(\frac{(\frac{1}{\pi} \langle \alpha G_{\mu\nu}^2 \rangle / F_\pi^4)}{(\frac{1}{\pi} \langle \alpha G_{\mu\nu}^2 \rangle / f_\pi^4)_{\text{QCD}}} \right)^{1/4} = \left(\frac{(\frac{1}{\pi} \langle \alpha G_{\mu\nu}^2 \rangle / F_\pi^4)}{(0.012 \text{ GeV}^4 / (92.4 \text{ MeV})^4)_{\text{QCD}}} \right)^{1/4}. \quad (3.12)$$

In Fig. 2 we show the plots of (M_ρ/F_π) , (M_{a_1}/F_π) , and (M_{TD}/F_π) as a function of Γ for $\gamma_m \simeq 0$ (left panel) and $\gamma_m \simeq 1$ (right panel) with $N_{TC} = 3$ and $\hat{S} = 0.31$ (QCD value) fixed, where in calculating (M_{TD}/F_π) we have used $\kappa = 1.0$. It is interesting to note that, for both cases with $\gamma_m \simeq 0$ and $\gamma_m \simeq 1$, (M_ρ/F_π) and (M_{a_1}/F_π) tend to coincide to be degenerate constant ($\simeq 8.0$) as Γ increases, fairly independent of the value of the S parameter: Such a degenerate spectrum of the techni- ρ and techni- a_1 relatively independent of the S parameter is a salient feature of the large contribution of the techni-gluon condensate and would have characteristic impact on the techni-hadron phenomenology. On the other hand, the *techni-dilaton* mass (M_{TD}/F_π) is very sensitive to Γ , rapidly decreasing as Γ increases. We have checked that these behaviors do not alter even for different values of $\hat{S}$ other than $\hat{S} = 0.31$.

It is also worth studying the γ_m -dependences of (M_ρ/F_π) , (M_{a_1}/F_π) , and (M_{TD}/F_π) . Figure 3 shows the plots of (M_ρ/F_π) , (M_{a_1}/F_π) , and (M_{TD}/F_π) as a function of γ_m for $\hat{S} = 0.31$ (left panel) and $\hat{S} = 0.1$ (right panel) with $N_{TC} = 3$ and $\Gamma = 1$ fixed. Looking at Fig. 3, we see that (M_ρ/F_π) and (M_{a_1}/F_π) are not sensitive to γ_m , while (M_{TD}/F_π) rapidly decreases as γ_m becomes larger for both cases with $\hat{S} = 0.31$ and $\hat{S} = 0.1$. Actually, in the case of $\hat{S} \leq 0.1$, such a relatively large suppression on (M_{TD}/F_π) generated by the large anomalous dimension can be seen by analytically solving Eq.(2.51) for $(M_{TD}z_m) \ll 1$ ^{#6} expanding Eq.(2.51) in power of $(M_{TD}z_m)$:

$$(M_{TD}z_m)^2 \simeq (2 - \gamma_m) \cdot 4\lambda\xi^2 \frac{g_5^2}{L} = (2 - \gamma_m) \cdot 3\kappa\xi^2, \quad (3.13)$$

where we have used Eqs.(3.2) and (3.3).

Let us next focus on the case of a generic SWC-TC with $\gamma_m \simeq 1$ and evaluate contributions from the techni-gluon condensation to the S parameter ($S = \hat{S} \cdot (N_f/2)$). In Fig. 4 we show the plot of $\hat{S}$ divided by N_{TC} as a function of $(F_\pi/M_\rho)^2$ for $\Gamma = 0, 5, 10$. In this figure $(\hat{S}/N_{TC})$ is restricted in a phenomenologically arrowed range, $(\hat{S}/N_{TC}) \lesssim 0.05$ ^{#7}. Looking at Fig. 4, we can easily see that $\hat{S}$ monotonically decreases with respect to $(F_\pi/M_\rho)^2$ in

^{#6} The condition $(M_{TD}z_m) \ll 1$ is satisfied only when $\hat{S} < 0.1$.

^{#7} The value of the upper bound, 0.05, can be estimated as follows: Consider a conservative upper bound of the S parameter, $S \lesssim 0.1$ and look at the relationship with $\hat{S}$, $(S/N_{TC}) = (N_f/2) \cdot (\hat{S}/N_{TC})$. Since $N_{TC} \geq 2$ and $N_f \geq 2$ (i.e. the number of techni-doublets $N_{TD} \geq 1$), one then finds that $S \lesssim 0.1$ leads to $(\hat{S}/N_{TC}) \lesssim (0.1/[N_{TC}]_{\min}) \times (2/[N_f]_{\min}) = 0.05$.




FIG. 3: Plots of (M_ρ/F_π) , (M_{a_1}/F_π) , and (M_{TD}/F_π) as a function of γ_m with $N_{TC} = 3$ and $\Gamma = 1$ fixed. The left panel corresponds to the case with $\hat{S} = 0.31$ while the right panel $\hat{S} = 0.1$. The dotted (red), dashed (yellow), solid (blue) lines respectively denote (M_ρ/F_π) , (M_{a_1}/F_π) , and (M_{TD}/F_π) in both panels.




FIG. 4: Plots of $\hat{S}/N_{TC}$ as a function of $(F_\pi/M_\rho)^2$ in the case of a generic SWC-TC with $\gamma_m \simeq 1$. In the left panel $\hat{S}/N_{TC}$ is restricted in a phenomenologically interesting region $\hat{S}/N_{TC} \lesssim 0.05$. The right panel focuses on the dashed-rectangle area ($\hat{S}/N_{TC} \lesssim 0.01$) in the left panel. In both panels, the thick (black) line represents the case with $\Gamma = 0$ which corresponds to the previous analysis without G [39], while the thin (blue) line and the dashed (red) line respectively correspond to the cases with $\Gamma = 5$ and $\Gamma = 10$.

both cases with $\Gamma = 0$ and $\Gamma \neq 0$ and is continuously approaching zero as $(F_\pi/M_\rho)^2 \rightarrow 0$. Conversely, when we tune $\hat{S}$ smaller, (M_ρ/F_π) gets larger ((M_{a_1}/F_π) and (M_{TD}/F_π) as well):

$$(M_\rho/F_\pi, M_{a_1}/F_\pi, M_{TD}/F_\pi) \nearrow \text{ as } \hat{S} \searrow. \quad (3.14)$$

This tendency coincides with what the authors found in Ref. [39]. Indeed, the model with $\Gamma = 0$ is nothing but the one analyzed in Ref. [39]. Note that for $\Gamma = 5$ and 10 the techni-gluon condensation reduces the S parameter up till 10% in the phenomenologically acceptable region of S ($S = \hat{S} \cdot (N_f/2) \lesssim 0.1$).

To see a general tendency of (M_ρ/F_π) , (M_{a_1}/F_π) , and (M_{TD}/F_π) with respect to Γ , we shall fix $\hat{S}$ to be a certain value, say, $\hat{S} = 0.1$ (minimal requirement for a realistic TC) ^{#8}. For $\hat{S} = 0.1$ and $N_{TC} = 3$, we evaluate the Γ -dependences of (M_ρ/F_π) , (M_{a_1}/F_π) , and (M_{TD}/F_π) to get the plots shown in Fig. 5. Here we have taken $\kappa = 1$ in calculating M_{TD} and have chosen $\Gamma = 10$ as the maximal value which turns out to be large enough when we consider typical models of SWC-TC as will be seen later. Figure 5 tells us that, for $0 \leq \Gamma \leq 10$, the input $\hat{S} = 0.1$ constrains the meson masses to be respectively in the following ranges: $15.7 \geq (M_\rho/F_\pi) \geq 14.6$, $17.3 \geq (M_{a_1}/F_\pi) \geq 14.7$, $9.79 \geq (M_{TD}/F_\pi) \geq 1.45$. This implies that the gluonic-contribution decreases (M_ρ/F_π) , (M_{a_1}/F_π) , and (M_{TD}/F_π) about 9%, 15%, and 85%, respectively, during the value of Γ evolves from 0 to 10. Note that, although we have not specified any types of SWC-TC yet, it is remarkable that each value of the M_ρ and M_{a_1} is fairly insensitive to Γ but becomes a degenerate non-zero constant as Γ increases, the value being almost determined just by fixing the S parameter. (The degeneracy itself is realized no matter what value the S might take.) In contrast, the mass of techni-dilaton M_{TD} is

^{#8} Actually, the phenomenological bound for the S parameter is $S \lesssim 0.1$, not $\hat{S}(= S/(N_f/2)) \lesssim 0.1$. Both constraints would coincide only when $N_f = 2$ (minimal flavors). More detailed discussion including the flavor-dependence will be given in the next section.




FIG. 5: Plots of (M_ρ/F_π) , (M_{a_1}/F_π) , and (M_{TD}/F_π) as a function of Γ for $\hat{S} = 0.1$ in a generic SWC-TC with $N_{TC} = 3$ and $\gamma_m \simeq 1$. The dotted (red), dashed (yellow), and solid (blue) curves respectively correspond to (M_ρ/F_π) , (M_{a_1}/F_π) , and (M_{TD}/F_π) .

very sensitive to Γ , rapidly decreasing as Γ increases. In fact $M_{TD}/F_\pi \rightarrow 0$ in the formal limit $\Gamma \rightarrow \infty$. This tendency is still operative even if S is much smaller, although the decreasing rate of (M_{TD}/F_π) becomes somewhat milder. In the next section we will discuss a matching with a couple of concrete models of SWC-TC in which the value of Γ is related to (Λ_{ETC}/F_π) in such a way that $\Gamma \simeq 6 - 8$ for $(\Lambda_{ETC}/F_\pi) = 10^4 - 10^5$. ($\Gamma \rightarrow \infty$ for $\Lambda_{ETC}/F_\pi \rightarrow \infty$.)

IV. MATCHING WITH TYPICAL MODELS OF SWC-TC BASED ON LARGE N_f QCD

In this section, we consider typical models of SWC-TC based on the Caswell-Banks-Zaks IR fixed point [16, 17] α_* of the two-loop beta function in the large N_f QCD, QCD with massless flavors $3 \ll N_f < 11N_{TC}/2$. It has been suggested that there exists a certain region ($N_f^* < N_f^{\text{cr}} < N_f < 11N_{TC}/2$) (which is called “conformal window”) such that $\alpha_* < \alpha_{\text{cr}}$, where the critical coupling α_{cr} for the spontaneous chiral symmetry breaking may be estimated as $\alpha_{\text{cr}} = \pi/(3C_2(F))$ based on the Schwinger-Dyson (SD) equation in the ladder approximation. Equating $\alpha_* = \alpha_{\text{cr}} = \pi/(3C_2(F))$, we find $N_f^{\text{cr}} \simeq 4N_{TC}$ [14] for the conformal phase transition point. There are many studies about the existence of the conformal window and the value of N_f^{cr} (if exists) in various non-perturbative methods including lattice gauge theories [48]. Here we simply assume the existence of the conformal window and tentatively use the value of N_f^{cr} from the perturbative two-loop beta function and ladder SD equation.

It was argued [15] that the conformal phase transition is characterized by scaling of the essential singularity (Miranisky scaling [24])

$$m \simeq 4\Lambda \exp\left(-\frac{\pi}{\sqrt{\alpha/\alpha_{\text{cr}} - 1}}\right), \quad (4.1)$$

where m is the dynamical mass of techni-fermion and Λ an intrinsic scale (Λ_{TC}) which may be identified with an ETC scale Λ_{ETC} in the actual model building:

$$\Lambda = \Lambda_{TC} \simeq \Lambda_{ETC} \gg m. \quad (4.2)$$

We can arrange a large hierarchy $m \ll \Lambda$ in terms of (approximately) conformal symmetry by tuning the theory close to the conformal fixed point as $\alpha \simeq \alpha_* \rightarrow \alpha_{\text{cr}}$ in the broken phase $\alpha_* > \alpha_{\text{cr}}$, in such a way that the coupling constant is almost non-running over the wide range $m < \mu < \Lambda$ (see Fig. 1). (There still exists a remnant of the conformal symmetry due to the IR fixed point α_* , although the IR fixed point α_* actually disappears because techni-fermions with mass m decouple from the beta function for $\mu < m$.)

In the SWC dynamics near the conformal window, the *explicit* breaking of the conformal symmetry manifest as the conformal anomaly is due to the generation of the dynamical mass of techni-fermions m which arises from the *spontaneous* breaking of the conformal and chiral symmetry. Thus the techni-gluon condensation for the conformal anomaly relevant to the dynamics near the conformal window is directly related to m but not the intrinsic scale $\Lambda_{TC} \gg m$. The conformal anomaly for this dynamical generation takes the following form [22]:

$$\langle \partial^\mu D_\mu \rangle = \langle \theta_\mu^\mu \rangle = 4\langle \theta_0^0 \rangle = \lim_{\Lambda \rightarrow \infty} \frac{\beta_{\text{NP}}(\alpha)}{4\alpha^2} \langle \alpha G_{\mu\nu}^2 \rangle, \quad (4.3)$$

where D_μ and θ_μ^ν are the dilatation current and the energy-momentum tensor, respectively, and $\beta_{\text{NP}}(\alpha)$ denotes the *non-perturbative* beta function of the gauge coupling α related to the generation of m in Eq.(4.1):

$$\beta_{\text{NP}}(\alpha) \equiv \frac{\partial \alpha}{\partial \ln \Lambda} = -\frac{2}{3C_2(F)} \left(\frac{\alpha}{\alpha_{\text{cr}}} - 1 \right)^{\frac{3}{2}} = -\frac{2\pi^3}{3C_2(F)} \left(\ln 4 \frac{\Lambda}{m} \right)^{-3}, \quad (4.4)$$

in which we have used $\alpha_{\text{cr}} = \pi/(3C_2(F))$. Straightforward calculation of the effective potential at two-loop order [22] yields the vacuum energy $\langle \theta_0^0 \rangle$:

$$\langle \theta_0^0 \rangle = -\frac{N_f N_{\text{TC}}}{\pi^4} m^4. \quad (4.5)$$

From Eqs.(4.3) and (4.5), we obtain

$$\langle \alpha G_{\mu\nu}^2 \rangle = -\frac{16}{\pi^4} \frac{\alpha^2}{\beta_{\text{NP}}(\alpha)} N_f N_{\text{TC}} m^4. \quad (4.6)$$

Note that Eq.(4.3) takes the same form as a conventional conformal anomaly obtained in the all order perturbation theory [3], where we naturally expect $\langle \alpha G_{\mu\nu}^2 \rangle = \mathcal{O}(\Lambda_{\text{TC}}^4) \gg \mathcal{O}(m^4)$. Here we ignore techni-gluon dilaton with mass of this order Λ_{TC} associated with the perturbative anomaly and the running effect of the coupling for $\mu > \Lambda_{\text{TC}}$. In the ordinary QCD with $N_c = N_f = 3$, we have $\langle \alpha G_{\mu\nu}^2 \rangle = \mathcal{O}(\Lambda_{\text{QCD}}^4) = \mathcal{O}(m^4)$: There is no such a large hierarchy $m \ll \Lambda_{\text{QCD}}$ and conformal region $m < \mu < \Lambda_{\text{QCD}}$ where the coupling constant is almost non-running $\alpha(Q) \simeq \text{constant}$. In contrast, our techni-gluon condensate $\langle \alpha G_{\mu\nu}^2 \rangle$ in Eq.(4.3) is responsible for the conformal anomaly induced by the dynamical generation of mass m and hence $\langle \alpha G_{\mu\nu}^2 \rangle = \mathcal{O}(m^4) \ll \mathcal{O}(\Lambda_{\text{TC}}^4)$ and our techni-dilaton is a bound state of techni-fermions with mass m which breaks the conformal symmetry near the conformal window. Also note that the conformal region with the almost non-running coupling is realized only when we arrange N_f as $N_f \propto N_{\text{TC}}$ in such a way that the fermionic dynamics and the gluonic dynamics cooperate intimately. Thus in Eq.(4.6) we have $\langle G_{\mu\nu}^2 \rangle \sim \mathcal{O}(N_f N_{\text{TC}}) \sim \mathcal{O}(N_{\text{TC}}^2)$ in accord with large N_{TC} counting relevant to holographic models.

From Eqs.(4.1), (4.4), and (4.6), the techni-gluon condensate $\langle \alpha G_{\mu\nu}^2 \rangle$ is expressed in terms of m and Λ as

$$\langle \alpha G_{\mu\nu}^2 \rangle|_{\alpha \simeq \alpha_*} = \lim_{\Lambda \rightarrow \infty} \frac{8}{3} \frac{N_f N_{\text{TC}}}{C_2(F) \pi^2} m^4 \left(\frac{\ln \frac{4\Lambda}{m}}{\pi} \right)^3 \left(1 + \left(\frac{\ln \frac{4\Lambda}{m}}{\pi} \right)^{-2} \right)^2. \quad (4.7)$$

Comparing this to Eq.(2.26) with Eq.(3.2) taken into account, we arrive at a relationship between G , $(z_m m)$, and (Λ_{ETC}/m) ,

$$G = C \cdot (z_m m)^4 \left(\frac{\ln \frac{4\Lambda_{\text{ETC}}}{m}}{\pi} \right)^3 \left(1 + \left(\frac{\ln \frac{4\Lambda_{\text{ETC}}}{m}}{\pi} \right)^{-2} \right)^2, \quad (4.8)$$

where

$$C = \frac{1}{2\pi} \frac{N_f N_{\text{TC}}}{N_{\text{TC}}^2 - 1}. \quad (4.9)$$

In terms of m , the techni-fermion condensate $\langle \bar{T}T \rangle_m$ renormalized at $\mu = m$ can be evaluated as ^{#9}

$$\langle \bar{T}T \rangle_m = Z_m \cdot \langle \bar{T}T \rangle_\Lambda = -t \cdot \frac{N_{\text{TC}}}{4\pi^2} m^3 \quad \text{with } t \simeq 2, \quad (4.10)$$

where the mass renormalization constant is $Z_m = Z_m(m/\Lambda) = m/\Lambda$. We regard the condensate in Eq.(2.27) as the one renormalized at $\mu = (1/L)$ following the procedure suggested in Ref. [39]. We then find the parameter ξ is related to G together with $(z_m m)$ as follows:

$$\xi = \frac{\sqrt{3}}{1+G} (z_m m)^2, \quad (4.11)$$

^{#9} Numerically t coincides with the prefactor in Eq.(2.27), $3 - \gamma_m$, for $\gamma_m \simeq 0, 1, 2$. In the case of QCD with $\gamma_m \simeq 0$, this implies $m \simeq 453 \text{ MeV}$ for the value of $\langle \bar{q}q \rangle \simeq -(277 \text{ MeV})^3$ in accord with the conventional constituent quark mass $m \simeq 350 \text{ MeV}$ and with $m \simeq 420 \text{ GeV}$ from the Pagels-Stokar (PS) formula. For details see Appendix. B

N_{TC}	$\log_{10}(\Lambda_{ETC}/F_\pi)$	ξ	G	$1/z_m$ [TeV]
2	4	0.22	1.12	2.09
2	5	0.15	1.69	2.31
3	4	0.21	0.60	1.90
3	5	0.16	0.97	2.04
4	4	0.19	0.37	1.82
4	5	0.15	0.62	1.92

TABLE II: Values of the model-parameters fitted to the SWC-TC with $N_{TC} = 2, 3, 4$ and $N_f = N_f^{EW} = 4N_{TC}$ based on the large N_f QCD. Here use has been made of $S = \hat{S}(N_f^{EW}/2) = 0.1$.

N_{TC}	$\log_{10}(\Lambda_{ETC}/F_\pi)$	M_{TD} [GeV]	M_ρ [TeV]	M_{a_1} [TeV]	Γ	m [TeV]	m/m_{PS}	R
2	4	777 $\left(\begin{smallmatrix} +106 \\ -125 \end{smallmatrix}\right)$	3.75	3.82	5.93	1.08	1.49	1.11
2	5	613 $\left(\begin{smallmatrix} +85 \\ -99 \end{smallmatrix}\right)$	3.69	3.74	7.26	1.13	1.57	1.16
3	4	681 $\left(\begin{smallmatrix} +94 \\ -110 \end{smallmatrix}\right)$	3.86	3.90	6.26	0.84	1.74	1.48
3	5	556 $\left(\begin{smallmatrix} +77 \\ -90 \end{smallmatrix}\right)$	3.80	3.83	7.57	0.87	1.80	1.53
4	4	597 $\left(\begin{smallmatrix} +82 \\ -96 \end{smallmatrix}\right)$	3.93	3.95	6.58	0.71	1.97	1.84
4	5	505 $\left(\begin{smallmatrix} +70 \\ -82 \end{smallmatrix}\right)$	3.87	3.89	7.88	0.73	2.02	1.88

TABLE III: Estimates of M_ρ , M_{a_1} , and M_{TD} for $S = 0.1$ in the SWC-TC with $N_{TC} = 2, 3, 4$ and $N_f = N_f^{EW} = 4N_{TC}$ based on the large N_f QCD. The range of the values of M_{TD} come from varying the value of κ in the range $0.7 \leq \kappa \leq 1.3$, where the smallest values of M_{TD} correspond to the cases with $\kappa = 0.7$ while the largest values $\kappa = 1.3$. m_{PS} and R are defined in the text.

where we have used Eq.(3.2). From Eqs.(4.11) and (4.8), we see that the two parameters ξ and G are now replaced by $(z_m m)$ and (Λ_{ETC}/m) involving the quantities concerning the SWC-TC, the dynamical mass m and the ETC scale Λ_{ETC} .

To be concrete for our analysis, we will assume the value of N_f^{cr} as that from the two-loop beta function and ladder SD equation $N_f^{cr} \simeq 4N_{TC}$ bearing in mind the large N_{TC} limit in accord with holographic setup.

Let us now recall Eqs.(3.7)-(3.11) which imply that all the quantities given in those equations (say, $S = S(\xi, G)$ and $F_\pi = z_m^{-1} \tilde{f}_\pi(\xi, G)$) are determined once we fix the one dimensionful parameter z_m and the two dimensionless parameters (ξ, G) which are now rephrased by $(z_m m)$ and (Λ_{ETC}/m) . To fix ξ or G , we may use a certain value of the S parameter (e.g. $S = 0.1$) as a phenomenological input in such a way as was done in the previous section. To determine the size of z_m , we can use the familiar formula

$$F_\pi = 246 / \sqrt{N_f^{EW}/2} \text{ GeV}, \quad (4.12)$$

where N_f^{EW} is the number of techni-fermions belonging to the doublet of $SU(2)_L$ symmetry in the standard model (SM) and may be different from the number of techni-fermion flavors N_f which participate in the SWC-TC dynamics. Furthermore, the ETC scale Λ_{ETC} may be constrained to be in a range, $10^3 \lesssim \Lambda_{ETC} \lesssim 10^4$ TeV:

$$10^4 \lesssim \Lambda_{ETC}/F_\pi \lesssim 10^5, \quad (4.13)$$

so as to accommodate the realistic light quark masses without suffering from the flavor changing neutral current (FCNC) syndrome. Use of these inputs now fixes the values of the parameters $(\xi, G, 1/z_m)$. In Table II we list these values for each $N_{TC} = 2, 3, 4$, and $(\Lambda_{ETC}/F_\pi) = 10^4, 10^5$ in the case of $N_f = N_f^{EW} = 4N_{TC}$ and $S = \hat{S}(N_f^{EW}/2) = 0.1$.

Using the values of the parameters given in Table II, we calculate the masses of the techni- ρ (M_ρ), techni- a_1 (M_{a_1}), and techni-scalar (M_{TD}) mesons to obtain Table III. The values of M_{TD} are estimated by varying the value of κ from 0.7 to 1.3 (around 1.0 with 30% error) in Table III. Note that M_ρ and M_{a_1} are almost degenerate to be $M_\rho \simeq M_{a_1} \simeq 3.7\text{--}3.9$ TeV for every $N_{TC} = 2, 3, 4$, in accord with the general tendency of the model with large techni-gluon condensate in section III where the degeneracy was not linked to the smallness of the S parameter and would have a new phenomenological implications. In contrast, the techni-dilaton mass $M_{TD} \sim 500\text{--}800$ GeV (when

$\kappa = 1$) is much lighter than M_ρ and M_{a_1} also in accord with the generic analysis for large Γ in the previous section, although $\hat{S} = S/(N_f^{\text{EW}}/2) = 0.1/(2N_{\text{TC}})$ is somewhat smaller than $\hat{S} = 0.1$ used in the generic analysis in Fig. 5. Indeed, in the present case we have $\Gamma \simeq 6-8$ as is seen from Table III. The essential reason for the large Γ is due to the existence of the wide conformal region $F_\pi < \mu < \Lambda_{\text{ETC}}$ with $(\Lambda_{\text{ETC}}/F_\pi) = 10^4-10^5$, which yields the smallness of the beta function through the factor $(\ln 4\Lambda_{\text{ETC}}/m)^{-3}$ in Eq.(4.4) and hence amplifies the techni-gluon condensation in Eq.(4.6) compared with the ordinary QCD with $\Gamma = 1$. Note that in the idealized (phenomenologically uninteresting) limit $\Lambda_{\text{ETC}}/F_\pi \rightarrow \infty$ we would have $\Gamma \rightarrow \infty$ and hence $M_{\text{TD}}/F_\pi \rightarrow 0$. (This does not mean that techni-dilaton becomes a true (exactly massless) NG boson, since its decay constant diverges, $F_{\text{TD}}/F_\pi \rightarrow \infty$, in such an idealized limit and hence the techni-dilaton gets decoupled. See later discussions.)

Thus we would expect the techni-dilaton as a composite Higgs boson near the conformality of SWC-TC with mass

$$M_{\text{TD}} \simeq 600 \text{ GeV}, \quad (4.14)$$

while M_ρ and M_{a_1} are generally heavy:

$$M_\rho \simeq M_{a_1} \simeq 3.8 \text{ TeV}. \quad (4.15)$$

The values of the ratio (m/m_{PS}) are also listed in Table. III, where m_{PS} denotes the dynamical mass estimated based on the PS formula:

$$F_\pi^2 \simeq \frac{N_{\text{TC}}}{4\pi^2} m_{\text{PS}}^2 \cdot I, \quad \text{with } I = \int_0^\infty dx x \frac{\Sigma^2(x) - \frac{x}{4} \frac{d}{dx} \Sigma^2(x)}{(x + \Sigma^2(x))^2}, \quad (4.16)$$

in which $\Sigma(x) \equiv \Sigma(Q^2)/m$ with $\Sigma(Q^2)$ being the mass function. When we use a simple-minded parametrization for $\Sigma(x)$, $\Sigma(x) = x^{(\gamma_m-1)/2}$ for $x = Q^2/m^2 > 1$ and $\Sigma(x) = 1$ for $x < 1$, we get $I \simeq 1$ for SWC-TC with $\gamma_m \simeq 1$ while $I \simeq 0.6$ for QCD with $\gamma_m \simeq 0$ (For details see Appendix. B). It is interesting to note that $m/m_{\text{PS}} > 1$. This can be understood by considering a PS formula appropriate for the present holographic analysis [39]:

$$F_\pi^2 \simeq \frac{N_{\text{TC}}}{4\pi^2} m^2 \cdot (z_m m)^{4-2\gamma_m}, \quad (4.17)$$

which is satisfied for $(1/z_m) > m$ and $0 \lesssim \gamma_m \lesssim 1$. Using Eqs.(4.16) and (4.17), certainly we find that $(m/m_{\text{PS}}) \simeq (\frac{1/z_m}{m}) > 1$ for $\gamma_m \simeq 1$.

Also listed is the ratio R defined as

$$R^3 = \sqrt{\frac{N_{\text{TC}}}{3}} \frac{\langle \bar{T}T \rangle_m / F_\pi^3}{(\langle \bar{q}q \rangle / f_\pi^3)_{\text{QCD}}}, \quad (4.18)$$

which is slightly larger than 1 reflected by the result that $m/m_{\text{PS}} > 1$, while it would be smaller than 1 ($R \simeq 0.69$) if the PS formula were used. Note that the enhancement of R has nothing to do with that of $\langle \bar{T}T \rangle_\Lambda$ caused by $Z_m^{-1}(\Lambda/m)$.

Let us now compare the present holographic model having the explicit techni-gluon contribution with the previous model [39] without the techni gluon condensation. Were it not for the matching with the ladder SD analysis (namely without using Eqs.(4.1),(4.6), and (4.10)), the model of Ref. [39] would be simply the $\Gamma = 0$ limit of the present model. Actually, the values of the meson masses lying on the $\Gamma = 0$ line in Fig. 5 are the results for $\hat{S} = 0.1$ in the previous model of Ref. [39]. Here we compare the two models for the same value of S as in Table. III. In Table IV we show the values of the meson masses for $S = 0.1$ and $N_{\text{TC}} = 2, 3, 4$ in the holographic SWC-TC without techni-gluon condensation [39], where we have used $\kappa = 1.0$ in estimating the values of M_{TD} . Comparing the values in Tables III (present model with $\Gamma \simeq 6-8$) and IV (previous model), one hardly sees differences in M_ρ and M_{a_1} , in accord with the general analysis in the previous section (see Fig. 2), while there is a substantial difference in M_{TD} arising from the non-zero techni-gluon condensation as seen in Fig. 2. Note that the relatively smaller M_{TD} compared with M_ρ and M_{a_1} is due to the large anomalous dimension as was mentioned in Eq.(3.13) and also seen from Fig. 3.

To summarize the characteristic feature of our holographic model, we may compare our result for $(M_\rho/F_\pi, M_{a_1}/F_\pi, M_{\text{TD}}/F_\pi)$ of the SWC-TC in Table III with that of the ordinary QCD ($N_c = N_f = 3$) in Table I. See Table.V where we show the case of $N_{\text{TC}} = 3$ and $(\Lambda_{\text{ETC}}/F_\pi) = 10^4$ as a representative case of our SWC-TC model (the first row) and the ordinary QCD with the input value of (M_ρ/f_π) (the second row), together with the $S = 0.1$ case of the previous holographic SWC-TC model without techni-gluon condensation ($\Gamma = 0$) [39] (the third row). Table V shows that the values of (M_ρ/F_π) and (M_{a_1}/F_π) tend to become larger as the value of S gets smaller as noted in the generic analysis (See Eq.(3.14)). This tendency is actually almost independent of the values of Γ and γ_m as seen in Figs. 2 and 3. Notably, although (M_{TD}/F_π) has the same tendency with respect to S , it receives other

N_{TC}	M_{TD} [TeV]	M_ρ [TeV]	M_{a_1} [TeV]
2	1.58	3.99	4.13
3	1.08	4.03	4.09
4	0.81	4.04	4.08

TABLE IV: Estimates of the meson mass in the holographic SWC-TC without techni-gluon condensation [39] for $S = 0.1$ with $N_{\text{TC}} = 2, 3, 4$ and $N_f = 4N_{\text{TC}}$. In calculating the values of M_{TD} the value of $\kappa = 1.0$ has been used.

Holographic models with $N_{\text{TC}}(\text{or } N_c) = 3$	M_ρ/F_π	M_{a_1}/F_π	M_{TD}/F_π
SWC-TC with $\gamma_m \simeq 1$ ($\Gamma = 7.57, S = 0.1, \Lambda_{\text{ETC}}/F_\pi = 10^5$)	37.8	38.1	5.5
QCD with $\gamma_m \simeq 0$ ($\Gamma = 1, S = 0.31$)	8.4	13.7	12.4
SWC-TC with $\gamma_m \simeq 1$ ($\Gamma = 0, S = 0.1$)	40.1	40.7	10.7

TABLE V: Comparison of $(M_\rho/F_\pi, M_{a_1}/F_\pi, M_{\text{TD}}/F_\pi)$ between several holographic models with $N_{\text{TC}}(\text{or } N_c) = 3$. In calculating the values of (M_{TD}/F_π) $\kappa = 1$ has been used.

important effects: One is from the large anomalous dimension $\gamma_m \simeq 1$ as in Fig. 3 and Eq.(3.13), while the other from the large techni-gluon condensation $\Gamma = 7.57$ as one can see from Figs. 2 and 5.

It is also worth comparing with the result of the straightforward calculation of the large N_f QCD with $N_{\text{TC}} = 3$ based on the Bethe-Salpeter (BS) equation combined with the SD equation in the ladder approximation [13, 23, 27]. The straightforward calculation of current correlators using the SD and BS equations has shown [13] that $\hat{S} \simeq 0.17$ ^{#10} near the conformal phase transition point, for $\alpha_*/\alpha_{\text{cr}} \simeq 1.13$, or equivalently, $(\Lambda_{\text{ETC}}/F_\pi) = (\Lambda/F_\pi) \simeq 10^{4.6}$. To make a direct comparison, we take the same value of $\hat{S}$ and $(\Lambda_{\text{ETC}}/F_\pi) = (\Lambda/F_\pi) \simeq 10^{4.6}$ ($\Gamma \simeq 5.09$). The comparison of the meson masses normalized by F_π is given in Table. VI. Here we have also shown the values with those in the previous holographic SWC-TC with $N_{\text{TC}} = 3$ without gluonic-condensation ($\Gamma = 0$) [39]. Looking at Table VI, one can see a curious coincidence among the values of (M_ρ/F_π) and (M_{a_1}/F_π) from different cases. This agreement does not depend so much on whether gluonic-contribution is turned off or not, as expected from Fig. 2. On the other hand, (M_{TD}/F_π) is somewhat smaller than the value calculated from the ladder BS equation with the SD equation.

SWC-TC models with $\gamma_m \simeq 1, N_{\text{TC}} = 3$ and $\hat{S} = 0.17$ [13]	M_ρ/F_π	M_{a_1}/F_π	M_{TD}/F_π
Holographic SWC-TC with $\Gamma = 5.09$ ($\Lambda_{\text{ETC}}/F_\pi = 10^{4.6}$)	11.2	11.5	2.3
Ladder BS with SD [23, 27]	11.0	11.5	4.2
Holographic SWC-TC with $\Gamma = 0$	11.6	13.9	8.9

TABLE VI: Comparison of the values of the meson masses normalized by F_π between three SWC-TC models with $\gamma_m \simeq 1$ and $N_{\text{TC}} = 3$. In calculating the values of (M_{TD}/F_π) in the holographic models $\kappa = 1$ has been used. F_π in Ref. [23, 27] is the value from the PS formula.

So far our analysis has been done for a specific value of S , $S = 0.1$. As we already noted in Eq.(3.14), $(M_\rho/F_\pi, M_{a_1}/F_\pi, M_{\text{TD}}/F_\pi)$ increase as S decreases. Here we show how our result changes for different values of S ($S \leq 0.1$). In Fig. 6 we show the values of M_ρ and M_{TD} for $\kappa = 1$ in the SWC-TC with $N_{\text{TC}} = 3$ ($N_f = 4N_{\text{TC}} = 12$). The shaded area is the region obtained for $10^4 \leq (\Lambda_{\text{ETC}}/F_\pi) \leq 10^5$, so as to satisfy the phenomenological bound from the FCNC and $S \leq 0.1$. Figure 6 indicates that, even when $0.07 \lesssim S \leq 0.1$, M_{TD} can still lie in a region less than 1 TeV which is still in the discovery region at the Large Hadron Collider (LHC).

Finally, we calculate the decay constant of techni-dilaton, F_{TD} , following a hypothesis of the partially conserved

^{#10} In Ref. [13] the straightforward calculation of current correlators in the large N_f QCD actually gives $\hat{S} \simeq 0.25$ for $N_{\text{TC}} = 3$. However, since it is known that the ladder SD and BS method has a tendency to overestimate $\hat{S}$ in QCD, which could be understood as scale ambiguity, the actual value near the conformal phase transition point with $\gamma_m \simeq 1$ has to be re-scaled by a factor about $(2/3)$ so as to fit the QCD value properly in extending to the case of QCD. Then we obtain the re-scaled value of $\hat{S}$, $\hat{S}^{\text{re-scaled}} \simeq 0.17$.




FIG. 6: Phenomenological constraints on masses of the techni- ρ and techni-dilaton in the SWC-TC model with $N_{\text{TC}} = 3$ ($N_f = 4N_{\text{TC}} = 12$) based on the large N_f QCD. The shaded region is drawn by varying the value of $(\Lambda_{\text{ETC}}/F_\pi)$ from 10^4 to 10^5 so as to satisfy the phenomenological constraint $S \leq 0.1$ and the FCNC. The dashed lines correspond to $S = 0.1, 0.09, 0.08, 0.07$ from bottom to top. In the plot the value of κ is taken to be 1.

dilatation current (PCDC) ^{#11} together with using the formula of the conformal anomaly given in Eq.(4.3). In this framework, the decay constant F_{TD} is given by [49]

$$F_{\text{TD}}^2 = -4 \frac{\langle \theta_\mu^\mu \rangle}{M_{\text{TD}}^2} = \frac{16N_f N_{\text{TC}}}{\pi^4} \frac{m^4}{M_{\text{TD}}^2}, \quad (4.19)$$

where we have used Eq.(4.5). This implies that $F_{\text{TD}}/F_\pi \sim m/M_{\text{TD}} \rightarrow \infty$ (decoupled techni-dilaton !) in the idealized (phenomenologically uninteresting) limit $\Gamma \rightarrow \infty$ ($\Lambda_{\text{ETC}}/F_\pi \rightarrow \infty$) where the techni-dilaton would be a true (exactly massless) Nambu-Goldstone boson, $M_\rho/F_\pi \sim M_{\text{TD}}/m \rightarrow 0$. In our framework the conformal symmetry is always broken explicitly as well as spontaneously. Table VII shows a list of the values of F_{TD} for each $N_{\text{TC}} = 2, 3, 4$ with $N_f = 4N_{\text{TC}}$, $(\Lambda_{\text{ETC}}/F_\pi) = 10^4$, and $\hat{S} = 0.1$ fixed. The value of F_{TD} may be compared with that obtained based on the linear σ model (LSM) approach which gives a formula between F_{TD} and F_π as $F_{\text{TD}}^{\text{LSM}} = (3 - \gamma_m)\sqrt{N_f/2}F_\pi = (3 - \gamma_m) \cdot 246 \text{ GeV} = 2 \times 246 \text{ GeV}$ (the third column of Table VII). The discrepancy between F_{TD} and $F_{\text{TD}}^{\text{LSM}}$ is not surprising since the SWC-TC dynamics is near the conformal phase transition where the Ginzburg-Landau/Gell-Mann-Levy effective theory (linear σ model) breaks down [15].

We can estimate the Yukawa coupling constant of techni-dilaton by F_{TD} as

$$g_Y^{\text{TD}} = \sqrt{2}(3 - \gamma_m)m_f/F_{\text{TD}}, \quad (4.20)$$

which is smaller than the Yukawa coupling in the SM, $g_Y^{\text{SM}} = \sqrt{2}m_f/(246 \text{ GeV}) = \sqrt{2}(3 - \gamma_m)m_f/F_{\text{TD}}^{\text{LSM}}$, as $g_Y^{\text{TD}}/g_Y^{\text{SM}} = (F_{\text{TD}}/F_{\text{TD}}^{\text{LSM}})^{-1} \simeq 0.18\text{--}0.20$. Such a difference may be useful for experimental search for the techni-dilaton in LHC.

N_{TC}	F_{TD} [TeV]	$F_{\text{TD}}/F_{\text{TD}}^{\text{LSM}}$	$g_Y^{\text{TD}}/g_Y^{\text{SM}}$
2	2.42	4.91	0.20
3	2.49	5.07	0.20
4	2.71	5.52	0.18

TABLE VII: Estimates of the values of F_{TD} for each $N_{\text{TC}} = 2, 3, 4$ with $(\Lambda_{\text{ETC}}/F_\pi) = 10^4$ and $\hat{S} = 0.1$ fixed. The values of the ratios $F_{\text{TD}}/F_{\text{TD}}^{\text{LSM}}$ and $g_Y^{\text{TD}}/g_Y^{\text{SM}}$ are also displayed.

^{#11} We could calculate F_{TD} , not invoking the PCDC hypothesis, straightforwardly from the scalar-current correlator based on the holographic principle. Such an alternative (but direct) calculation would be pursued in future publications.

V. SUMMARY AND DISCUSSIONS

In this paper, we have proposed a holographic SWC-TC including the bulk flavor/chiral-singlet scalar field Φ_X corresponding to (techni-) gluon condensation, based on deformation of a hard-wall-type bottom-up holographic QCD by adjusting the anomalous dimension γ_m . Thanks to the additional explicit bulk scalar field Φ_X , we naturally reproduced gluonic $1/Q^4$ terms in the OPE of the underlying theory (QCD and SWC-TC) for current correlators, in such a way as clearly to distinguish them from the same $1/Q^4$ terms due to the chiral condensate in the case of SWC-TC with $\gamma_m \simeq 1$. We have analyzed a generic case with $0 \lesssim \gamma_m \lesssim 1$ and calculated the masses of the techni- ρ meson (M_ρ), the techni- a_1 meson (M_{a_1}), and the flavor-singlet scalar meson, techni-dilaton (M_{TD}), as well as the S parameter.

We have shown that our model with $\gamma_m = 0$ and $N_f = 3$ well reproduces the real-life QCD (See Table I), with an indication that the QCD ρ meson mass M_ρ includes a (negative) contribution about 10% from the gluon condensate (See Eq.(3.6)).

For the case of $\gamma_m = 1$, we studied the dependence of the S parameter on (M_ρ/F_π) for several values of the techni-gluon condensation Γ (Eq.(3.12)), $\Gamma = 0, 5, 10$ (Fig. 4): $\hat{S}$ decreases monotonically with respect to (F_π/M_ρ) to continuously approach zero. This implies (M_ρ/F_π) necessarily increases when $\hat{S}$ is required to be smaller (Eq.(3.14)). It was also shown that, in the region of $\hat{S} \lesssim 0.1$, the techni-gluon condensation reduces the value of S maximally about 10%, compared with the previous analysis without techni-gluon contribution [39].

In the generic TC case with $0 \lesssim \gamma_m \lesssim 1$, we discussed how the meson masses relative to F_π , $(M_\rho/F_\pi, M_{a_1}/F_\pi, M_{TD}/F_\pi)$ change by varying γ_m , $\hat{S}$, and the techni-gluon condensation Γ . For fixed value of $\hat{S}(=0.31)$ (QCD value, Fig. 2) and $\hat{S} = 0.1$ (minimal requirement for a realistic TC, Fig. 5), (M_ρ/F_π) and (M_{a_1}/F_π) are sensitive to neither Γ nor γ_m , although the degeneracy between (M_ρ/F_π) and (M_{a_1}/F_π) takes place for somewhat larger Γ . In contrast, the techni-dilaton mass has a characteristic feature related to the conformality of SWC-TC: (M_{TD}/F_π) substantially decreases as Γ and/or γ_m increases when $\hat{S}$ is fixed. Particularly, $M_{TD}/F_\pi \rightarrow 0$ in the formal limit $\Gamma \rightarrow \infty$.

To specify the value of Γ , we considered a couple of typical models of SWC-TC with $\gamma_m \simeq 1$ and $N_{TC} = 2, 3, 4$ based on the Caswell-Banks-Zaks infrared fixed point in the large N_f QCD. Using some specific dynamical features of the conformal anomaly indicated by the analysis based on the ladder SD equation, we found the relation of Γ to (Λ_{ETC}/F_π) : In the case of $N_{TC} = 3$ ($N_f = 4N_{TC}$) and $S = (N_f/2) \cdot \hat{S} \simeq 0.1$, we have $\Gamma \simeq 7$ for $(\Lambda_{ETC}/F_\pi) = 10^4\text{--}10^5$ (required by the FCNC constraint). Thanks to the large anomalous dimension γ_m and large techni-gluon condensation Γ , we had a relatively light techni-dilaton with mass $M_{TD} \simeq 600$ GeV compared with $M_\rho \simeq M_{a_1} \simeq 3.8$ TeV (Table III). Note that large values of M_ρ and M_{a_1} are essentially determined by the requirement of $S = 0.1$ fairly independently of the techni-gluon condensation Γ , though the degeneracy between them is due to the largeness of Γ . Such large values of M_ρ and M_{a_1} might make the standard signatures through these techni-hadrons quite invisible at LHC. Note however that the largeness of the spectra in our model simply comes from the constraint from the S parameter *evaluated by the TC sector alone*. If we found other effects to reduce the S parameter such as the ETC effects implementing the mass of the SM fermions, we could pull down the overall scale of the whole techni-hadron spectra, in which case the degenerate techni- ρ and techni- a_1 as well as the lighter techni-dilaton in our model would have much impact in the LHC phenomenology.

The essential reason for the large Γ is due to the existence of the wide conformal region $F_\pi < \mu < \Lambda_{ETC}$ with $(\Lambda_{ETC}/F_\pi) = 10^4\text{--}10^5$, which yields the smallness of the beta function through the factor $(\ln 4\Lambda_{ETC}/m)^{-3}$ (see Eq.(4.4)) and hence amplifies the techni-gluon condensation in Eq.(4.6) compared with the ordinary QCD with $\Gamma = 1$. In the idealized (phenomenologically uninteresting) limit $\Lambda_{ETC}/F_\pi \rightarrow \infty$ we would have $\Gamma \rightarrow \infty$ and hence $M_{TD}/F_\pi \rightarrow 0$. The would-be “massless” techni-dilaton is actually decoupled, since its decay constant F_{TD} becomes divergent in that limit as is implied in Eq.(4.19). There exists no isolated massless techni-dilaton in contrast to the chiral symmetry breaking: The scale symmetry is broken both spontaneously and explicitly.

The predicted mass of holographic techni-dilaton (“conformal Higgs”), 600 GeV, lies in the discovery region at LHC. The size of Yukawa coupling of the techni-dilaton was also estimated through the PCDC relation, which turned out to be somewhat smaller than that of the SM Higgs (Table. VII). More detailed analysis on intrinsic signals at LHC concerning such a holographic techni-dilaton/conformal Higgs will be explored in future publications.

Before closing this section, several comments are in order:

A. Perturbative unitarity-bound and partially EW gauged SWC-TC

Since the vector meson masses predicted from our analysis are of order of a few TeV and hence irrelevant to the unitarity (See Table III), it would be the light techni-dilaton that is responsible for the perturbative unitarity of $W_L W_L$ scattering. The perturbative unitarity-bound on M_{TD} can then be estimated through a formula,

$$M_{\text{TD}} \lesssim \Lambda_{\text{uni}} = \sqrt{8\pi} F_\pi = \sqrt{8\pi} \cdot \left(\frac{246 \text{ GeV}}{\sqrt{N_f^{\text{EW}}/2}} \right), \quad (5.1)$$

where N_f^{EW} denotes the number of the techni-fermions charged under the electroweak (EW) gauge. In the case of the SWC-TC models with $N_f^{\text{EW}} = N_f = 4N_{\text{TC}}$ listed in Table III, however, the unitarity bound on the values of M_{TD} may be estimated as

$$M_{\text{TD}} \Big|_{\text{Table III}} \lesssim (617, 503, 436) \text{ GeV}, \quad \text{for } N_{\text{TC}} = 2, 3, 4. \quad (5.2)$$

Looking at the values of M_{TD} listed in Table III, we see that, for every case of N_{TC} , some masses of the techni-dilaton are somewhat heavier than that required by the unitarity bound above.

The situation with the perturbative unitarity would be much improved in a class of models (so-called partially gauged model) considered for other purpose [50]: Only a part of techni-fermion flavors carries the EW charges, while other fermion flavors are EW-singlets introduced only to achieve the SWC behavior of technicolor dynamics. Here the idea is to relax our condition $N_f^{\text{EW}} = N_f = 4N_{\text{TC}}$ such that $N_f^{\text{EW}} < N_f (= 4N_{\text{TC}}) = N_f^{\text{EW}} + N_f^{\text{EW-singlet}}$, which increases $\hat{S} = S \cdot (2/N_f^{\text{EW}})$ for the same $S (= 0.1)$ compared with the analysis in Sec. IV. Actually, $M_{\text{TD}}/\Lambda_{\text{uni}} = \frac{1}{\sqrt{8}}(M_{\text{TD}}/F_\pi)$ decreases when $\hat{S}$ increases as shown in Figs. 3, 4, and Eq.(3.14). In Table VIII we give two examples of the partially gauged models with $N_f = 4N_{\text{TC}}$ (one-doublet models with $N_f^{\text{EW}} = 2$ and one-family models with $N_f^{\text{EW}} = 8$), where the unitarity condition is fulfilled, $M_{\text{TD}}/\Lambda_{\text{uni}} \lesssim 1$.

one-family partially gauged model $N_f^{\text{EW}} = 8 \leq 4N_{\text{TC}}$ ($\Lambda_{\text{uni}} = \sqrt{8\pi} F_\pi \simeq 617 \text{ GeV}$)				
N_{TC}	$\log_{10}(\Lambda_{\text{ETC}}/F_\pi)$	M_{TD} [GeV]	M_ρ [TeV]	M_{a_1} [TeV]
2	4	777 $\left(\begin{smallmatrix} +106 \\ -125 \end{smallmatrix}\right)$	3.75	3.82
2	5	613 $\left(\begin{smallmatrix} +85 \\ -99 \end{smallmatrix}\right)$	3.69	3.74
3	4	725 $\left(\begin{smallmatrix} +99 \\ -117 \end{smallmatrix}\right)$	3.81	3.86
3	5	581 $\left(\begin{smallmatrix} +81 \\ -94 \end{smallmatrix}\right)$	3.74	3.78
4	4	686 $\left(\begin{smallmatrix} +94 \\ -110 \end{smallmatrix}\right)$	3.84	3.88
4	5	557 $\left(\begin{smallmatrix} +77 \\ -90 \end{smallmatrix}\right)$	3.78	3.81
one-doublet partially gauged model with $N_f^{\text{EW}} = 2 < 4N_{\text{TC}}$ ($\Lambda_{\text{uni}} = \sqrt{8\pi} F_\pi \simeq 1.23 \text{ TeV}$)				
N_{TC}	$\log_{10}(\Lambda_{\text{ETC}}/F_\pi)$	M_{TD} [GeV]	M_ρ [TeV]	M_{a_1} [TeV]
2	4	863 $\left(\begin{smallmatrix} +119 \\ -140 \end{smallmatrix}\right)$	3.57	3.70
2	5	676 $\left(\begin{smallmatrix} +94 \\ -110 \end{smallmatrix}\right)$	3.57	3.65
3	4	821 $\left(\begin{smallmatrix} +113 \\ -133 \end{smallmatrix}\right)$	3.61	3.71
3	5	645 $\left(\begin{smallmatrix} +90 \\ -105 \end{smallmatrix}\right)$	3.59	3.71
4	4	792 $\left(\begin{smallmatrix} +109 \\ -128 \end{smallmatrix}\right)$	3.65	3.66
4	5	623 $\left(\begin{smallmatrix} +87 \\ -101 \end{smallmatrix}\right)$	3.62	3.68

TABLE VIII: Estimates of the meson masses in one-family and one-doublet partially gauged models based on the large N_f QCD with $S = 0.1$ fixed. The values of M_{TD} are estimated varying the value of κ in the range $0.7 \leq \kappa \leq 1.3$, where the smallest values of M_{TD} correspond to the cases with $\kappa = 0.7$, while the largest values $\kappa = 1.3$.




FIG. 7: The plot of $\hat{S}/N_{TC}$ as a function of $\log_{10}[\Lambda/m]$ for $M_\rho/m = 2$.

B. Possible scaling behaviors of S and the conformal phase transition/chiral restoration

The analyses in this paper were made for fixed values of $\hat{S}$ or S ($\neq 0$) and $m/\Lambda_{TC} \simeq F_\pi/\Lambda_{TC} = 10^{-4}-10^{-5}$. The simplest extension of our analysis would imply that $S \sim (F_\pi/M_\rho)^2 \rightarrow \text{constant} \neq 0$ as $m/\Lambda_{TC} \rightarrow 0$, which, as indicated in Table VI, seems to be in accord with the straightforward calculation based on ladder SD and BS equation [13]: $\hat{S} \simeq \text{constant} \simeq 0.17$ near $\alpha_*/\alpha_{cr} = 1.13$ ($F_\pi/\Lambda_{TC} = 10^{-4.6}$) and $\hat{S} \rightarrow \text{constant} \neq 0$ in the extrapolation toward $\alpha_*/\alpha_{cr} \rightarrow 1$ ($F_\pi/\Lambda_{TC} \rightarrow 0$).

Here we shall mention different possibilities for the scaling of S , other than $\hat{S} \rightarrow \text{constant}$, arising as $m/\Lambda_{TC} \rightarrow 0$ near the conformal phase transition. In the case of conformal phase transition, all the dimensionful parameters of order $\mathcal{O}(m) \ll \mathcal{O}(\Lambda_{TC})$ are expected to become zero at the critical point, $m/\Lambda_{TC} \rightarrow 0$. Therefore, vector meson mass also goes to zero, $M_\rho/\Lambda_{TC} \rightarrow 0$. Suppose first that $M_\rho/m = \text{constant}$ in the limit $m/\Lambda_{TC} \rightarrow 0$. Dimensionless parameter S is written in terms of two independent dimensionless parameters, say, ξ and G , once we fix γ_m . A set of ξ and G is converted to another set of dimensionless parameters M_ρ/m and Λ/m through Eqs.(4.8) and (4.11). If we fix $M_\rho/m = \text{constant}$, then S is given as a function of Λ/m . In Fig. 7, we show the plot of $\hat{S}/N_{TC}$ as a function of Λ/m for $\gamma_m = 1$ and $M_\rho/m = 2$. Actually, it turns out that $\hat{S}$ goes to zero with the chiral restoration, i.e., $\Lambda/m \rightarrow \infty$, whatever the constant value of M_ρ/m is taken to be. (If M_ρ were taken to be bigger than $2m$, $\hat{S}$ would become smaller.) This would imply $F_\pi/m \sim F_\pi/M_\rho \rightarrow 0$ and hence $\langle \bar{T}T \rangle_m/F_\pi^3 \sim m^3/F_\pi^3 \rightarrow \infty$ at the conformal phase transition point. This is the behavior somewhat different from that expected from the PS formula [39].

Acknowledgments

We would like to thank M. Hashimoto and M. Harada for useful comments and fruitful discussions. This work was supported in part by the Grant-in-Aid for Nagoya University Global COE Program, “Quest for Fundamental Principles in the Universe: from Particles to the Solar System and the Cosmos”, from the Ministry of Education, Culture, Sports, Science and Technology of Japan. It was also supported by the JSPS Grant-in-Aid for Scientific Research (S) #22224003. S.M. was supported by the Korean Research Foundation Grant funded by the Korean Government (KRF-2008-341-C00008).

Appendix A: Holographic matching to the operator product expansion of current correlators

We start with Eq.(2.30), the equation of motion for $V_\mu(q, z)$, and rewrite it in terms of $V(q, z)$ defined as $V_\mu(q, z) = \tilde{v}_\mu(q)V(q, z)$ as follows:

$$[Q^2 + w^{-1}(z)\partial_z w(z)\partial_z]V(q, z) = 0, \quad (\text{A.1})$$

where $Q^2 = -q^2$ denotes a Euclidean momentum-squared and the induced metric $w(z)$ is given (in the limit $M' = 0$) as

$$w(z) = \frac{L}{z} \left(1 + G \left(\frac{z}{z_m} \right)^4 \right)^2. \quad (\text{A.2})$$

From Eq.(2.26) we note that

$$\frac{G}{z_m^4 Q^4} = -\frac{1}{8} \frac{c g_5^4}{L^2} \frac{\langle \alpha G_{\mu\nu}^2 \rangle}{Q^4}. \quad (\text{A.3})$$

Using this relation, we express the induced metric $w(z)$ as

$$w(z = y/Q) = \frac{QL}{y} \left(1 - \frac{1}{8} \frac{c g_5^4}{L^2} y^4 \frac{\langle \alpha G_{\mu\nu}^2 \rangle}{Q^4} \right)^2, \quad (\text{A.4})$$

where the variable z has been replaced with $y = Qz$.

Consider a large Euclidean momentum region $(1/z_m)^2 \ll Q^2 < (1/\epsilon)^2$. Expanding Eq.(A.1) in powers of Q^2 and using Eq.(2.35), we then obtain an asymptotic form of the vector current correlator $\Pi_V(Q^2)$,

$$\Pi_V(Q^2) \Big|_{(1/z_m)^2 \ll Q^2 < (1/\epsilon)^2} = Q^2 \left[\frac{L}{2g_5^2} \ln Q^2 + c \frac{2}{3} \frac{g_5^2}{L} \frac{\langle \alpha G_{\mu\nu}^2 \rangle}{Q^4} + \mathcal{O}\left(\frac{1}{Q^8}\right) \right]. \quad (\text{A.5})$$

This expression may be compared with the form of the operator product expansion (OPE) for arbitrary γ_m ,

$$\Pi_V(Q^2) \Big|_{\text{OPE}} = Q^2 \left[\frac{N_{\text{TC}}}{24\pi^2} \log \left(\frac{Q^2}{\mu^2} \right) - \frac{1}{24\pi} \frac{\langle \alpha G_{\mu\nu}^2 \rangle}{Q^4} + 2\pi \frac{\alpha \langle \bar{T}T \rangle_{\mu}^2 \mu^{-2\gamma_m}}{Q^{2(3-\gamma_m)}} \right]. \quad (\text{A.6})$$

Then we find the following matching conditions:

$$\frac{L}{g_5^2} = \frac{N_{\text{TC}}}{12\pi^2}, \quad c = -\frac{1}{16\pi} \frac{L}{g_5^2} = -\frac{N_{\text{TC}}}{192\pi^3}. \quad (\text{A.7})$$

Similar discussion on the axial-vector current correlator Π_A provides the same matching conditions for (L/g_5^2) and c as it should.

The high-energy expansion form in Eq.(A.5) does not include the chiral condensate term behaving as $1/Q^{2(3-\gamma_m)}$ as in Eq.(A.6). In order to incorporate this missing term, we shall introduce higher dimensional interaction terms constructed from the bulk scalar field Φ and the left- and the right-gauge fields L_M and R_M .

We consider the following dimension-six operators which are invariant under the five-dimensional $SU(N_f)_L \times SU(N_f)_R$ gauge symmetry:

$$\Delta\mathcal{L}_5 = \frac{L^2}{g_5^2} e^{c g_5^2 \Phi_X} \left(\frac{C_{LL}}{2} \text{Tr} [\Phi^\dagger \Phi] \text{Tr} [L_{MN} L^{MN} + R_{MN} R^{MN}] + C_{LR} \text{Tr} [\Phi^\dagger L_{MN} \Phi R^{MN}] \right), \quad (\text{A.8})$$

where C_{LL} and C_{LR} are the dimensionless coupling constants. The Lagrangian $\Delta\mathcal{L}_5$ gives shifts to the kinetic terms of the vector ($V_M \equiv (L_M + R_M)/\sqrt{2}$) and axial-vector ($A_M \equiv (L_M - R_M)/\sqrt{2}$) gauge fields as follows:

$$\frac{L^2}{g_5^2} e^{c g_5^2 \Phi_X} \left(\frac{C_V}{2} v^2(z) \text{Tr} [V_{MN} V^{MN}] + \frac{C_A}{2} v^2(z) \text{Tr} [A_{MN} A^{MN}] \right), \quad (\text{A.9})$$

where $C_{V,A} = C_{LL} \pm \frac{1}{2} C_{LR}$.

Let us focus on the vector sector taking $C_A = 0$ for simplicity. Then we see that the C_V term in Eq.(A.9) modifies the induced metric $w(z)$ in Eq.(A.2) as

$$w(z) \rightarrow \tilde{w}(z) = \frac{L}{z} \left(1 + G \left(\frac{z}{z_m} \right)^4 \right)^2 (1 - C_V L^2 v^2(z)). \quad (\text{A.10})$$

It should be noted from Eqs.(2.27) and (2.22)-(2.24) that $v^2(z)$ in Eq.(A.10) is expressed (in the chiral limit $M = 0$) as

$$v^2(z = y/Q) \simeq \frac{y^{2(3-\gamma_m)}}{3(3-\gamma_m)^2} \left(\frac{g_5^2}{L} \right)^2 \frac{\langle \bar{T}T \rangle_{1/L}^2 L^{2\gamma_m-2}}{Q^{2(3-\gamma_m)}}, \quad (\text{A.11})$$

where we have neglected higher order terms in $1/Q^2$ expansion. Thus we see that the modified-induced metric $\tilde{w}(z)$ now includes the desired $\bar{T}T$ term:

$$\tilde{w}(z = y/Q) \ni \frac{QL}{y} \left[-C_V \left(\frac{g_5^2}{L} \right)^2 \frac{y^{2(3-\gamma_m)}}{3(3-\gamma_m)^2} \frac{L^{2\gamma_m} \langle \bar{T}T \rangle_{1/L}^2}{Q^{2(3-\gamma_m)}} \right]. \quad (\text{A.12})$$

Combining this with Eq.(A.4), we obtain the total expression of the high-energy expansion for Π_V :

$$\Pi_V(Q^2) \Big|_{(1/z_m)^2 \ll Q^2 < (1/\epsilon)^2} = Q^2 \left[\frac{L}{2g_5^2} \ln Q^2 + c \frac{2}{3} \frac{g_5^2}{L} \frac{\langle \alpha G_{\mu\nu}^2 \rangle}{Q^4} + C_6^V L^{2\gamma_m} \frac{\langle \bar{T}T \rangle_{1/L}^2}{Q^{2(3-\gamma_m)}} + \mathcal{O}\left(\frac{1}{Q^8}\right) \right], \quad (\text{A.13})$$

where

$$C_6^V = C_V \frac{\sqrt{\pi}}{6} \frac{(\Gamma(2-\gamma_m))^3}{\Gamma(\frac{7}{2}-\gamma_m)} \frac{g_5^2}{L}. \quad (\text{A.14})$$

Comparing Eq.(A.13) with the $\langle \bar{T}T \rangle^2$ term in Eq.(A.6), we determine the coefficient C_V as

$$C_V = 12\sqrt{\pi}\alpha \frac{\Gamma(\frac{7}{2}-\gamma_m)}{(\Gamma(3-\gamma_m))^3} \cdot \frac{L}{g_5^2}. \quad (\text{A.15})$$

For $C_A \neq 0$, one can similarly perform the high-energy expansion of Π_A . As a consequence of matching with the OPE for Π_A , we get

$$C_A = -12\sqrt{\pi}\alpha \frac{\Gamma(\frac{7}{2}-\gamma_m)}{(\Gamma(3-\gamma_m))^3} \frac{L}{g_5^2} + \frac{1}{(3-\gamma_m)(2-\gamma_m)}. \quad (\text{A.16})$$

Thus, it is shown that the present model with the higher dimensional terms in Eq.(A.8) added completely reproduces the OPE for the current correlators up to terms suppressed by $1/Q^8$.

Appendix B: The Pagels-Stokar formula and chiral condensate

1. Relationship between decay constant and dynamical fermion mass

The Pagels-Stokar formula relates the mass function $\Sigma(Q^2)$ with F_π as

$$F_\pi^2 = \frac{N_{\text{TC}}}{4\pi^2} m_{\text{PS}}^2 \int_0^{(\Lambda^2/m_{\text{PS}}^2 \rightarrow \infty)} dx x \frac{\Sigma^2(x) - \frac{x}{4} \frac{d}{dx} \Sigma^2(x)}{(x + \Sigma^2(x))^2}, \quad (\text{B.1})$$

where $\Sigma(x) = \Sigma(Q^2)/m_{\text{PS}}$ which may be parametrized as

$$\Sigma(x) = \begin{cases} x^{\frac{\gamma_m}{2}-1} & \text{for } x > 1 \\ 1 & \text{for } x < 1 \end{cases}. \quad (\text{B.2})$$

By using Eq.(B.2), Eq.(B.1) is calculated as ^{#12}

$$\frac{4\pi^2}{N_{\text{TC}}} \frac{F_\pi^2}{m_{\text{PS}}^2} \simeq \left(\int_0^1 dx \frac{x}{(x+1)^2} + \frac{(3-\frac{\gamma_m}{2})}{2} \int_1^\infty dx \frac{x^{\gamma_m-1}}{(x+x^{\gamma_m-2})^2} \right)$$

^{#12} One may neglect $\Sigma^2(x) = x^{\gamma_m-2}$ in the denominator of the integrand of Eq.(B.1) since in the integration dominant contributions come from the UV region where $\Sigma^2(x) = x^{\gamma_m-2} \ll x$ for $0 \lesssim \gamma_m \lesssim 1$. Then the form of Eq.(B.3) would be changed to $\frac{4\pi^2}{N_{\text{TC}}} \frac{F_\pi^2}{m_{\text{PS}}^2} \simeq -\frac{1}{2} + \ln 2 + \frac{1}{2} \frac{(3-\frac{\gamma_m}{2})}{(2-\gamma_m)}$.

$$= -\frac{1}{2} + \ln 2 + \frac{1}{2(3-\gamma_m)^2} \left((3-\gamma_m) + H_{\frac{-1}{2(3-\gamma)}} - H_{\frac{-(4-\gamma)}{2(3-\gamma)}} \right), \quad (\text{B.3})$$

where H_n is harmonic number with order of n defined as $H_n = \sum_{k=1}^n 1/k^n$. For $\gamma_m = 0, 1$, Eq.(B.3) is evaluated as

$$\frac{4\pi^2}{N_{\text{TC}}} \frac{F_\pi^2}{m_{\text{PS}}^2} \simeq \begin{cases} 0.63 & \text{for } \gamma_m = 0 \\ 1.00 & \text{for } \gamma_m = 1 \end{cases}, \quad (\text{B.4})$$

and for $\gamma_m = 2$ we have $\frac{4\pi^2}{N_{\text{TC}}} \frac{F_\pi^2}{m_{\text{PS}}^2} \simeq \ln(\Lambda/m)^2$.

2. Relationship between chiral condensate and dynamical fermion mass

The chiral condensate $\langle \bar{T}T \rangle$ evaluated at a UV scale Λ is given as

$$\langle \bar{T}T \rangle_\Lambda = \frac{N_{\text{TC}}}{4\pi^2} m^3 \int_0^{(\Lambda^2/m^2 \rightarrow \infty)} dx \frac{x\Sigma(x)}{x + \Sigma^2(x)}. \quad (\text{B.5})$$

One can calculate Eq.(B.5) using the following relation derived based on analysis of the ladder SD equation [51] ^{#13}

$$\int_0^{(\Lambda^2/m^2 \rightarrow \infty)} dx \frac{x\Sigma(x)}{x + \Sigma^2(x)} = \left. \frac{\frac{d}{dx}\Sigma(x)}{\frac{d}{dx}(\lambda(x)/x)} \right|_{x=(\Lambda/m)^2}, \quad (\text{B.7})$$

where $\lambda(x) = \frac{1}{4} \frac{\alpha}{\alpha_{\text{cr}}}$ with $\alpha_{\text{cr}} = \pi/(3C_2(F))$.

In the case of SWC-TC with $\gamma_m \simeq 1$ where $\alpha(\simeq \alpha_*) \simeq \alpha_{\text{cr}}$, we have $\lambda = 1/4$. Using Eqs.(B.2) and (B.7) we then evaluate Eq.(B.5) as

$$-\frac{4\pi^2}{N_{\text{TC}}} \frac{\langle \bar{T}T \rangle_\Lambda}{m^3} \Big|_{\text{SWC-TC}} = 2 \cdot Z_m^{-1}, \quad (\text{B.8})$$

where $Z_m = (\Lambda/m)^{-1}$ is the mass renormalization constant.

In the case of QCD with $\gamma_m \simeq 0$, $\Sigma(x)$ and $\lambda(x)$ are expressed as

$$\Sigma(x) = \frac{1}{x} \left[1 + \frac{1}{2A} \ln x \right]^{A/2-1}, \quad (\text{B.9})$$

$$\lambda(x) = \frac{A/2}{\ln x + 2A}, \quad (\text{B.10})$$

where $A = 1/(b\alpha_{\text{cr}})$ with b being the coefficient of QCD beta function at one-loop order, $b = \frac{1}{6\pi} (11N_c - 2N_f)$. Substituting Eqs.(B.9) and (B.10) into Eq.(B.7), we have

$$\begin{aligned} -\frac{4\pi^2}{N_c} \frac{\langle \bar{T}T \rangle_\Lambda}{m^3} \Big|_{\text{QCD}} &\simeq \frac{4}{(2A)^{A/2}} \cdot Z_m^{-1} \\ &\simeq Z_m^{-1} \cdot \begin{cases} 3.1 & \text{for real life QCD with } N_c = N_f = 3 (A = 8/9) \\ 3.3 & \text{for large } N_c \text{ QCD } (A = 9/11) \end{cases}, \end{aligned} \quad (\text{B.11})$$

^{#13} Neglecting the $\Sigma(x)$ in the denominator, which is not dominant in the integral, and substituting Eq.(B.2) into Eq.(B.5), we may evaluate Eq.(B.5) to reach a form,

$$-\frac{4\pi^2}{N_{\text{TC}}} \frac{\langle \bar{T}T \rangle_\Lambda}{m^3} \simeq \left[1 - \ln 2 + \frac{2}{\gamma_m} \left(\left(\frac{\Lambda}{m} \right)^{\gamma_m} - 1 \right) \right], \quad (\text{B.6})$$

which turns out to be good approximation and gives the same numbers as those in Eqs.(B.8) and (B.12), except for the QCD case with $\gamma_m = 0$.

where $Z_m = (\ln(\Lambda^2/m^2))^{-A/2}$.

For the case of constant mass $\Sigma(x) = 1$ which corresponds to the case with $\gamma_m = 2$ as in the Nambu-Jona-Lasinio (NJL) model, we can straightforwardly calculate Eq.(B.5) to get

$$-\frac{4\pi^2}{N_{\text{TC}}} \frac{\langle \bar{T}T \rangle_\Lambda}{m^3} \Big|_{\text{NJL}} = 1 \cdot Z_m^{-1}, \quad (\text{B.12})$$

where $Z_m = (\Lambda^2/m^2)^{-1}$

Finally using $\langle \bar{T}T \rangle_m = Z_m \langle \bar{T}T \rangle_\Lambda$, we thus reach a result

$$-\frac{4\pi^2}{N_{\text{TC}}} \frac{\langle \bar{T}T \rangle_m}{m^3} \simeq \begin{cases} 3 & \text{for } \gamma_m = 0 \\ 2 & \text{for } \gamma_m = 1 \\ 1 & \text{for } \gamma_m = 2 \end{cases}. \quad (\text{B.13})$$

-
- [1] K. Haba, S. Matsuzaki and K. Yamawaki, arXiv:1003.2841 [hep-ph]. Talk given at Nagoya Global COE workshop: “Strong Coupling Gauge Theories in LHC Era” in honor of Toshihide Maskawa’s 70th Birthday and 35th Anniversary of Dynamical Symmetry Breaking in SCGT (SCGT 09), Dec. 8-11, 2009, Nagoya University, Nagoya. To be published by World Scientific Publishing Co., Singapore (eds. M. Harada, H. Fukaya, M. Tanabashi and K. Yamawaki).
 - [2] S. Weinberg, Phys. Rev. D **13**, 974 (1976); L. Susskind, Phys. Rev. D **20**, 2619 (1979).
 - [3] S. L. Adler, J. C. Collins and A. Duncan, Phys. Rev. D **15**, 1712 (1977); J. C. Collins, A. Duncan and S. D. Joglekar, Phys. Rev. D **16**, 438 (1977); N. K. Nielsen, Nucl. Phys. B **120**, 212 (1977).
 - [4] M. E. Peskin and T. Takeuchi, Phys. Rev. Lett. **65**, 964 (1990); Phys. Rev. D **46**, 381 (1992); B. Holdom and J. Terning, Phys. Lett. B **247**, 88 (1990); M. Golden and L. Randall, Nucl. Phys. B **361**, 3 (1991).
 - [5] K. Yamawaki, M. Bando and K. Matumoto, Phys. Rev. Lett. **56**, 1335 (1986); M. Bando, K. Matumoto and K. Yamawaki, Phys. Lett. B **178**, 308 (1986); M. Bando, T. Morozumi, H. So and K. Yamawaki, Phys. Rev. Lett. **59**, 389 (1987).
 - [6] T. Akiba and T. Yanagida, Phys. Lett. B **169**, 432 (1986).
 - [7] T. W. Appelquist, D. Karabali and L. C. R. Wijewardhana, Phys. Rev. Lett. **57**, 957 (1986); T. Appelquist and L. C. R. Wijewardhana, Phys. Rev. D **36**, 568 (1987).
 - [8] B. Holdom, Phys. Lett. B **150**, 301 (1985).
 - [9] T. Maskawa and H. Nakajima, Prog. Theor. Phys. **52**, 1326 (1974).
 - [10] V. A. Miransky, *Dynamical Symmetry Breaking in Quantum Field Theories* (World Scientific Pub. Co., Singapore 1993); K. Yamawaki, hep-ph/9603293, in *Proceedings of 14th Symposium on Theoretical Physics “Dynamical Symmetry Breaking and Effective Field Theory”*, Cheju Island, Korea, July 21-26, 1995, ed. J. E. Kim (Minumsa Pub. Co., Korea, 1996) p.43-86; C. T. Hill, and E. H. Simmons, Phys. Rept. **381** (2003), 235 [Erratum *ibid.* **390** (2004), 553].
 - [11] S. Dimopoulos and L. Susskind, Nucl. Phys. B **155**, 237 (1979); E. Eichten and K. D. Lane, Phys. Lett. B **90**, 125 (1980).
 - [12] T. Appelquist and G. Triantaphyllou, Phys. Lett. B **278**, 345 (1992); R. Sundrum and S. D. H. Hsu, Nucl. Phys. B **391**, 127 (1993); T. Appelquist and F. Sannino, Phys. Rev. D **59**, 067702 (1999).
 - [13] M. Harada, M. Kurachi and K. Yamawaki, Prog. Theor. Phys. **115**, 765 (2006); M. Kurachi and R. Shrock, Phys. Rev. D **74**, 056003 (2006); M. Kurachi, R. Shrock and K. Yamawaki, Phys. Rev. D **76**, 035003 (2007).
 - [14] T. Appelquist, J. Terning and L. C. R. Wijewardhana, Phys. Rev. Lett. **77**, 1214 (1996); **79**, 2767 (1997); T. Appelquist, A. Ratnaweera, J. Terning and L. C. R. Wijewardhana, Phys. Rev. D **58**, 105017 (1998).
 - [15] V. A. Miransky and K. Yamawaki, Phys. Rev. D **55**, 5051 (1997) [Erratum-*ibid.* D **56**, 3768 (1997)].
 - [16] W. E. Caswell, Phys. Rev. Lett. **33**, 244 (1974).
 - [17] T. Banks and A. Zaks, Nucl. Phys. B **196**, 189 (1982).
 - [18] R. Fukuda and T. Kugo, Nucl. Phys. B **117** (1976), 250.
 - [19] Y. Iwasaki, K. Kanaya, S. Kaya, S. Sakai and T. Yoshie, Phys. Rev. D **69**, 014507 (2004).
 - [20] T. Appelquist, G. T. Fleming and E. T. Neil, Phys. Rev. Lett. **100**, 171607 (2008) [Erratum-*ibid.* **102**, 149902 (2009)]; A. Deuzeman, M. P. Lombardo and E. Pallante, Phys. Lett. B **670**, 41 (2008).
 - [21] F. Sannino and K. Tuominen, Phys. Rev. D **71**, 051901 (2005); D. K. Hong, S. D. H. Hsu and F. Sannino, Phys. Lett. B **597**, 89 (2004); For a review F. Sannino, arXiv:0804.0182 [hep-ph].
 - [22] V. A. Miransky and V. P. Gusynin, Prog. Theor. Phys. **81** (1989) 426.
 - [23] M. Harada, M. Kurachi and K. Yamawaki, Phys. Rev. D **68**, 076001 (2003).
 - [24] V. A. Miransky, Nuovo Cim. A **90**, 149 (1985).
 - [25] S. Shuto, M. Tanabashi and K. Yamawaki, in *Proc. 1989 Workshop on Dynamical Symmetry Breaking*, Dec. 21-23, 1989, Nagoya, eds. T. Muta and K. Yamawaki (Nagoya Univ., Nagoya, 1990) 115-123.
 - [26] M. S. Carena and C. E. M. Wagner, Phys. Lett. B **285**, 277 (1992); M. Hashimoto, Phys. Lett. B **441**, 389 (1998).
 - [27] M. Kurachi and R. Shrock, JHEP **0612**, 034 (2006).
 - [28] J. M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998) [Int. J. Theor. Phys. **38**, 1113 (1999)].
 - [29] N. Arkani-Hamed, M. Porrati and L. Randall, JHEP **0108**, 017 (2001).

- [30] L. Da Rold and A. Pomarol, Nucl. Phys. B **721**, 79 (2005).
- [31] J. Erlich, E. Katz, D. T. Son and M. A. Stephanov, Phys. Rev. Lett. **95**, 261602 (2005).
- [32] T. Sakai and S. Sugimoto, Prog. Theor. Phys. **113**, 843 (2005).
- [33] M. Bando, T. Kugo, S. Uehara, K. Yamawaki and T. Yanagida, Phys. Rev. Lett. **54**, 1215 (1985); M. Bando, T. Kugo and K. Yamawaki, Nucl. Phys. B **259**, 493 (1985); Phys. Rept. **164**, 217 (1988); M. Bando, T. Fujiwara and K. Yamawaki, Prog. Theor. Phys. **79**, 1140 (1988).
- [34] M. Harada and K. Yamawaki, Phys. Rept. **381**, 1 (2003).
- [35] H. Georgi, Nucl. Phys. B **266**, 274 (1986); N. Arkani-Hamed, A. G. Cohen and H. Georgi, Phys. Rev. Lett. **86**, 4757 (2001).
- [36] K. Agashe, C. Csaki, C. Grojean and M. Reece, JHEP **0712**, 003 (2007).
- [37] D. K. Hong and H. U. Yee, Phys. Rev. D **74**, 015011 (2006).
- [38] M. Piai, arXiv:hep-ph/0608241.
- [39] K. Haba, S. Matsuzaki and K. Yamawaki, Prog. Theor. Phys. **120**, 691 (2008).
- [40] A. Karch, E. Katz, D. T. Son and M. A. Stephanov, Phys. Rev. D **74**, 015005 (2006).
- [41] L. Da Rold and A. Pomarol, JHEP **0601**, 157 (2006).
- [42] J. Hirn, N. Rius and V. Sanz, Phys. Rev. D **73**, 085005 (2006); J. Hirn and V. Sanz, Phys. Rev. Lett. **97**, 121803 (2006).
- [43] G. 't Hooft, G. Isidori, L. Maiani, A. D. Polosa and V. Riquer, Phys. Lett. B **662**, 424 (2008).
- [44] D. Black, A. H. Fariborz and J. Schechter, Phys. Rev. D **61**, 074001 (2000).
- [45] C. Amsler *et al.* [Particle Data Group], Phys. Lett. B **667**, 1 (2008).
- [46] M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **147**, 385 (1979); M. A. Shifman, A. I. Vainshtein and V. I. Zakharov, Nucl. Phys. B **147**, 448 (1979).
- [47] J. Gasser and H. Leutwyler, Phys. Rept. **87**, 77 (1982).
- [48] See for example talks in Nagoya SCGT 09 workshop. <http://www.eken.phys.nagoya-u.ac.jp/scgt09/>.
- [49] M. Bando, K. i. Matumoto and K. Yamawaki, Phys. Lett. B **178**, 308 (1986).
- [50] D. D. Dietrich, F. Sannino and K. Tuominen, Phys. Rev. D **72**, 055001 (2005); N. D. Christensen and R. Shrock, Phys. Lett. B **632**, 92 (2006).
- [51] M. Bando, T. Morozumi, H. So and K. Yamawaki, Phys. Rev. Lett. **59**, 389 (1987).